

# Coronavirus (COVID-19) - Diagnosed or Suspected

After Hours Telephone Triage Protocols | Adult | 2020

## DEFINITION

- Diagnosis was confirmed by **positive lab test** OR
- **Clinical diagnosis** (suspected diagnosis) was made by healthcare provider (HCP) OR
- **Patient or caregiver suspects COVID-19** based on symptoms consistent with COVID-19 and recalls close contact with a person with COVID-19 in past two weeks, or is living in an area of community spread.

### Note to Triager:

- Triagers should use their clinical judgment, but generally will want to use the *COVID-19 Diagnosed or Suspected* guideline when a patient calls with cough, fever, shortness of breath, or a combination of typical COVID symptoms and there is community spread.
- During the 2020-2021 influenza season, triagers can use the *COVID-19 Diagnosed or Suspected* guideline when a patient calls with flu-like symptoms.
- In adult patients triager should use clinical judgment to decide whether a symptom guideline (e.g., Chest Pain, Fever, Headache) should be used in addition to the *COVID-19 Diagnosed or Suspected* guideline. Adult patients have multiple factors that make assessment and triage more complex. Adults (and especially older adults) are more likely to have one or more chronic underlying medical conditions. Increased age is also a strong risk factor for severe COVID-19 illness and complications. Further, fever as a marker of important pathology increases significantly by age in adults.

This guideline was **last updated** 11/3/2020.

## INITIAL ASSESSMENT QUESTIONS

1. COVID-19 DIAGNOSIS: "Who made your Coronavirus (COVID-19) diagnosis?" "Was it confirmed by a positive lab test?" If not diagnosed by a HCP, ask "Are there lots of cases (community spread) where you live?" (See public health department website, if unsure)
2. COVID-19 EXPOSURE: "Was there any known exposure to COVID before the symptoms began?" CDC Definition of close contact: within 6 feet (2 meters) for a total of 15 minutes or more over a 24-hour period.
3. ONSET: "When did the COVID-19 symptoms start?"
4. WORST SYMPTOM: "What is your worst symptom?" (e.g., cough, fever, shortness of breath, muscle aches)
5. COUGH: "Do you have a cough?" If so, ask: "How bad is the cough?"
6. FEVER: "Do you have a fever?" If so, ask: "What is your temperature, how was it measured, and when did it start?"
7. RESPIRATORY STATUS: "Describe your breathing?" (e.g., shortness of breath, wheezing, unable to speak)
8. BETTER-SAME-WORSE: "Are you getting better, staying the same or getting worse compared to yesterday?" If getting worse, ask, "In what way?"
9. HIGH RISK DISEASE: "Do you have any chronic medical problems?" (e.g., asthma, heart or lung disease, weak immune system, obesity, etc.)
10. PREGNANCY: "Is there any chance you are pregnant?" "When was your last menstrual period?"
11. OTHER SYMPTOMS: "Do you have any other symptoms?" (e.g., chills, fatigue, headache, loss of smell or taste, muscle pain, sore throat; new loss of smell or taste especially support the diagnosis of COVID-19)

## TRIAGE ASSESSMENT QUESTIONS

### Call EMS 911 Now

SEVERE difficulty breathing (e.g., struggling for each breath, speaks in single words)

*R/O: respiratory failure, hypoxia*

- 1 Call EMS 911 Now:**
  - Immediate medical attention is needed. You need to hang up and call 911 (or an ambulance).
  - *Triager Discretion:* I'll call you back in a few minutes to be sure you were able to reach them.
- 2 Tell the Ambulance Dispatcher About COVID-19 Diagnosis:**
  - When you call 911, tell the dispatcher that you probably have COVID-19.
- 3 Tell Ambulance Medics About Your COVID-19 Diagnosis:**
  - Tell the paramedic right away that you probably have COVID-19.
  - The paramedics should call ahead to the emergency department to let them know.
- 4 Care Advice given per Coronavirus (COVID-19) - Diagnosed or Suspected (Adult) guideline.**

Difficult to awaken or acting confused (e.g., disoriented, slurred speech)

*R/O: hypoxia, sepsis*

- 1 Call EMS 911 Now:**
  - Immediate medical attention is needed. You need to hang up and call 911 (or an ambulance).
  - *Triager Discretion:* I'll call you back in a few minutes to be sure you were able to reach them.
- 2 Tell the Ambulance Dispatcher About COVID-19 Diagnosis:**
  - When you call 911, tell the dispatcher that you probably have COVID-19.
- 3 Tell Ambulance Medics About Your COVID-19 Diagnosis:**
  - Tell the paramedic right away that you probably have COVID-19.
  - The paramedics should call ahead to the emergency department to let them know.
- 4 Care Advice given per Coronavirus (COVID-19) - Diagnosed or Suspected (Adult) guideline.**

Bluish (or gray) lips or face now

*R/O: cyanosis and need for oxygen*

- 1 Call EMS 911 Now:**
  - Immediate medical attention is needed. You need to hang up and call 911 (or an ambulance).
  - *Triager Discretion:* I'll call you back in a few minutes to be sure you were able to reach them.
- 2 Tell the Ambulance Dispatcher About COVID-19 Diagnosis:**
  - When you call 911, tell the dispatcher that you probably have COVID-19.
- 3 Tell Ambulance Medics About Your COVID-19 Diagnosis:**
  - Tell the paramedic right away that you probably have COVID-19.
  - The paramedics should call ahead to the emergency department to let them know.
- 4 Care Advice given per Coronavirus (COVID-19) - Diagnosed or Suspected (Adult) guideline.**

Shock suspected (e.g., cold/pale/clammy skin, too weak to stand, low BP, rapid pulse)

*R/O: shock*

- 1 Call EMS 911 Now:**
  - Immediate medical attention is needed. You need to hang up and call 911 (or an ambulance).
  - *Triager Discretion:* I'll call you back in a few minutes to be sure you were able to reach them.
- 2 Tell the Ambulance Dispatcher About COVID-19 Diagnosis:**
  - When you call 911, tell the dispatcher that you probably have COVID-19.
- 3 Tell Ambulance Medics About Your COVID-19 Diagnosis:**
  - Tell the paramedic right away that you probably have COVID-19.
  - The paramedics should call ahead to the emergency department to let them know.
- 4 Care Advice given per Coronavirus (COVID-19) - Diagnosed or Suspected (Adult) guideline.**

Sounds like a life-threatening emergency to the triager

**1 Call EMS 911 Now:**

- Immediate medical attention is needed. You need to hang up and call 911 (or an ambulance).
- *Triager Discretion:* I'll call you back in a few minutes to be sure you were able to reach them.

**2 Tell the Ambulance Dispatcher About COVID-19 Diagnosis:**

- When you call 911, tell the dispatcher that you probably have COVID-19.

**3 Tell Ambulance Medics About Your COVID-19 Diagnosis:**

- Tell the paramedic right away that you probably have COVID-19.
- The paramedics should call ahead to the emergency department to let them know.

**4 Care Advice** given per **Coronavirus (COVID-19) - Diagnosed or Suspected (Adult)** guideline.

**See More Appropriate Guideline**

[1] COVID-19 exposure AND [2] no symptoms

*Go to Guideline: Coronavirus (COVID-19) - Exposure (Adult)*

[1] Lives with someone known to have influenza (flu test positive) AND [2] flu-like symptoms (e.g., cough, runny nose, sore throat, SOB; with or without fever)

*Go to Guideline: Influenza - Seasonal (Adult). Note: patient is more likely to have flu than COVID-19 if they are living with someone who tested positive for influenza.*

[1] Adult with possible COVID-19 symptoms AND [2] triager concerned about severity of symptoms or other causes

*Note to Triager: First use this guideline. In adult patients triager should then use clinical judgment to decide whether a symptom guideline (e.g., Chest Pain, Fever, Headache) should also be used.*

Immunization reaction suspected (e.g., fever, headache, muscle aches occurring during days 1-3 days after immunization)

*Go to Guideline: Immunization Reactions (Adult)*

COVID-19 and breastfeeding, questions about

*Go to Guideline: Coronavirus (COVID-19) - Diagnosed or Suspected (Pediatric)*

**Go to ED Now**

SEVERE or constant chest pain or pressure (Exception: mild central chest pain, present only when coughing)

*R/O: pneumonia, pleurisy. Note to Triager: Consider using both this guideline AND the Chest Pain guideline if any concern for cardiac or other more serious cause of chest pain.*

**1 Go to ED Now:**

- You need to be seen in the Emergency Department.
- Go to the ED at \_\_\_\_\_ Hospital.
- Leave now. Drive carefully.

- 2 **Note to Triager - Triage Nurse Should Notify Emergency Department (ED):**
  - Tell the ED you are sending a patient with suspected diagnosis of COVID-19 who is getting worse and inform them of patient's symptoms.
  - Obtain and document the patient / caller's mobile phone number. Either keep the patient on hold or call the patient back with instructions.
  - *Reason:* So that ED can make plans to prevent coronavirus spread to others in the hospital.
- 3 **Reassurance and Education - Going to the ED or Urgent Care Center During the COVID-19 Pandemic:**
  - If you or your child needs to be seen for an urgent medical problem, do not hesitate to go.
  - ERs and urgent care centers are safe places. They are well equipped to protect you against the virus.
  - For non-urgent conditions, talk to your healthcare provider's office first.
- 4 **You Should Go to the Emergency Department (ED):**
  - You will need to go to a nearby ED.
  - Do not leave until I've called and talked with the ED. The ED may have special instructions on how best to get you there. I will call you back (or place you on hold).
- 5 **You Should Tell Healthcare Personnel That You Might Have COVID-19:**
  - Tell the first healthcare worker you meet that you may have COVID-19.
  - Tell them you have symptoms and have been sent for COVID-19 testing.
- 6 **Wear a Mask - Cover Your Mouth and Nose:**
  - Wear a mask.
  - If you do not have a mask, then cover your mouth and nose with a disposable tissue (e.g., Kleenex, toilet paper, paper towel) or wash cloth.
- 7 **Another Adult Should Drive:**
  - It is better and safer if another adult drives instead of you.
- 8 **Call EMS 911 If:**
  - Severe difficulty breathing occurs
  - Lips or face turns blue
  - Confusion occurs.
- 9 **Care Advice given per Coronavirus (COVID-19) - Diagnosed or Suspected (Adult) guideline.**

MODERATE difficulty breathing (e.g., speaks in phrases, SOB even at rest, pulse 100-120)

*R/O: pneumonia*

- 1 **Go to ED Now:**
  - You need to be seen in the Emergency Department.
  - Go to the ED at \_\_\_\_\_ Hospital.
  - Leave now. Drive carefully.
- 2 **Note to Triager - Triage Nurse Should Notify Emergency Department (ED):**
  - Tell the ED you are sending a patient with suspected diagnosis of COVID-19 who is getting worse and inform them of patient's symptoms.
  - Obtain and document the patient / caller's mobile phone number. Either keep the patient on hold or call the patient back with instructions.
  - *Reason:* So that ED can make plans to prevent coronavirus spread to others in the hospital.
- 3 **Reassurance and Education - Going to the ED or Urgent Care Center During the COVID-19 Pandemic:**
  - If you or your child needs to be seen for an urgent medical problem, do not hesitate to go.
  - ERs and urgent care centers are safe places. They are well equipped to protect you against the virus.
  - For non-urgent conditions, talk to your healthcare provider's office first.
- 4 **You Should Go to the Emergency Department (ED):**
  - You will need to go to a nearby ED.
  - Do not leave until I've called and talked with the ED. The ED may have special instructions on how best to get you there. I will call you back (or place you on hold).
- 5 **You Should Tell Healthcare Personnel That You Might Have COVID-19:**
  - Tell the first healthcare worker you meet that you may have COVID-19.
  - Tell them you have symptoms and have been sent for COVID-19 testing.
- 6 **Wear a Mask - Cover Your Mouth and Nose:**
  - Wear a mask.
  - If you do not have a mask, then cover your mouth and nose with a disposable tissue (e.g., Kleenex, toilet paper, paper towel) or wash cloth.

- 7 **Another Adult Should Drive:**
  - It is better and safer if another adult drives instead of you.
- 8 **Call EMS 911 If:**
  - Severe difficulty breathing occurs
  - Lips or face turns blue
  - Confusion occurs.
- 9 **Care Advice** given per **Coronavirus (COVID-19) - Diagnosed or Suspected** (Adult) guideline.

[1] Headache AND [2] stiff neck (can't touch chin to chest)

*R/O: meningitis*

- 1 **Go to ED Now:**
  - You need to be seen in the Emergency Department.
  - Go to the ED at \_\_\_\_\_ Hospital.
  - Leave now. Drive carefully.
- 2 **Note to Triage - Triage Nurse Should Notify Emergency Department (ED):**
  - Tell the ED you are sending a patient with suspected diagnosis of COVID-19 who is getting worse and inform them of patient's symptoms.
  - Obtain and document the patient / caller's mobile phone number. Either keep the patient on hold or call the patient back with instructions.
  - *Reason:* So that ED can make plans to prevent coronavirus spread to others in the hospital.
- 3 **Reassurance and Education - Going to the ED or Urgent Care Center During the COVID-19 Pandemic:**
  - If you or your child needs to be seen for an urgent medical problem, do not hesitate to go.
  - ERs and urgent care centers are safe places. They are well equipped to protect you against the virus.
  - For non-urgent conditions, talk to your healthcare provider's office first.
- 4 **You Should Go to the Emergency Department (ED):**
  - You will need to go to a nearby ED.
  - Do not leave until I've called and talked with the ED. The ED may have special instructions on how best to get you there. I will call you back (or place you on hold).
- 5 **You Should Tell Healthcare Personnel That You Might Have COVID-19:**
  - Tell the first healthcare worker you meet that you may have COVID-19.
  - Tell them you have symptoms and have been sent for COVID-19 testing.
- 6 **Wear a Mask - Cover Your Mouth and Nose:**
  - Wear a mask.
  - If you do not have a mask, then cover your mouth and nose with a disposable tissue (e.g., Kleenex, toilet paper, paper towel) or wash cloth.
- 7 **Another Adult Should Drive:**
  - It is better and safer if another adult drives instead of you.
- 8 **Care Advice** given per **Coronavirus (COVID-19) - Diagnosed or Suspected** (Adult) guideline.

### Go to ED Now (or PCP triage)

MILD difficulty breathing (e.g., minimal/no SOB at rest, SOB with walking, pulse <100)

*R/O: pneumonia. Note: not from stuffy nose (e.g., not relieved by cleaning out the nose)*

- 1 **Go To ED Now (or PCP triage):**
  - **If No PCP (Primary Care Provider) Second-Level Triage:** You need to be seen within the next hour. Go to the ED/UCC at \_\_\_\_\_ Hospital. Leave as soon as you can.
  - **If PCP Second-Level Triage Required:** You may need to be seen. Your doctor (or NP/PA) will want to talk with you to decide what's best. I'll page the provider on-call now. If you haven't heard from the provider (or me) within 30 minutes, go directly to the ED/UCC at \_\_\_\_\_ Hospital.
- 2 **Alternate Disposition - Call Telemedicine Provider Now:**
  - Telemedicine may be your best choice for care during this COVID-19 outbreak.
  - You should call a telemedicine provider now, if your own healthcare provider is not available.

- 3 **Note to Triager - If NO PCP, Have Other HCP Re-triage the Patient, if Available:**
  - During this COVID-19 pandemic, the medical community is trying to prevent unnecessary referrals to the emergency department (ED). Some patients are fearful of being exposed to COVID-19 in a medical setting. Second-level triage (re-triage) by a physician has been shown to reduce ED referrals. Here are resources that may be available in your community.
  - **PCP Second-level Telephone Triage:** Some PCPs (primary care providers) want to provide re-triage before any of their non-emergent patients are referred to an ED. This requires their approval.
  - **Telemedicine:** Telemedicine is often a preferred source of second-level triage and care during this pandemic. Many practices and some hospitals now offer a telemedicine (virtual visit) service. There are also many national telemedicine companies that are delivering COVID-19 care.
  - **Emergency Department (ED):** Some EDs may provide a telephone follow-up service for patients who have COVID-19 with worsening symptoms.
- 4 **General Care Advice for COVID-19 Symptoms:**
  - The treatment is the same whether you have COVID-19, influenza or some other respiratory virus.
  - *Cough:* Use cough drops.
  - *Feeling dehydrated:* Drink extra liquids. If the air in your home is dry, use a humidifier.
  - *Fever:* For fever over 101 F (38.3 C), take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package. Do not take aspirin unless your doctor has prescribed it for you.
  - *Muscle aches, headache, and other pains:* Often this comes and goes with the fever. Take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package.
  - *Sore throat:* Try throat lozenges, hard candy or warm chicken broth.
- 5 **Care Advice** given per **Coronavirus (COVID-19) - Diagnosed or Suspected (Adult)** guideline.

## Chest pain or pressure

*R/O: pneumonia, pleurisy, chest discomfort from COVID-19*

- 1 **Go To ED Now (or PCP triage):**
  - **If No PCP (Primary Care Provider) Second-Level Triage:** You need to be seen within the next hour. Go to the ED/UCC at \_\_\_\_\_ Hospital. Leave as soon as you can.
  - **If PCP Second-Level Triage Required:** You may need to be seen. Your doctor (or NP/PA) will want to talk with you to decide what's best. I'll page the provider on-call now. If you haven't heard from the provider (or me) within 30 minutes, go directly to the ED/UCC at \_\_\_\_\_ Hospital.
- 2 **Alternate Disposition - Call Telemedicine Provider Now:**
  - Telemedicine may be your best choice for care during this COVID-19 outbreak.
  - You should call a telemedicine provider now, if your own healthcare provider is not available.
- 3 **Note to Triager - If NO PCP, Have Other HCP Re-triage the Patient, if Available:**
  - During this COVID-19 pandemic, the medical community is trying to prevent unnecessary referrals to the emergency department (ED). Some patients are fearful of being exposed to COVID-19 in a medical setting. Second-level triage (re-triage) by a physician has been shown to reduce ED referrals. Here are resources that may be available in your community.
  - **PCP Second-level Telephone Triage:** Some PCPs (primary care providers) want to provide re-triage before any of their non-emergent patients are referred to an ED. This requires their approval.
  - **Telemedicine:** Telemedicine is often a preferred source of second-level triage and care during this pandemic. Many practices and some hospitals now offer a telemedicine (virtual visit) service. There are also many national telemedicine companies that are delivering COVID-19 care.
  - **Emergency Department (ED):** Some EDs may provide a telephone follow-up service for patients who have COVID-19 with worsening symptoms.
- 4 **General Care Advice for COVID-19 Symptoms:**
  - The treatment is the same whether you have COVID-19, influenza or some other respiratory virus.
  - *Cough:* Use cough drops.
  - *Feeling dehydrated:* Drink extra liquids. If the air in your home is dry, use a humidifier.
  - *Fever:* For fever over 101 F (38.3 C), take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package. Do not take aspirin unless your doctor has prescribed it for you.
  - *Muscle aches, headache, and other pains:* Often this comes and goes with the fever. Take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package.
  - *Sore throat:* Try throat lozenges, hard candy or warm chicken broth.
- 5 **Care Advice** given per **Coronavirus (COVID-19) - Diagnosed or Suspected (Adult)** guideline.



## Patient sounds very sick or weak to the triager

*Reason: severe acute illness or serious complication suspected*

- 1 Go To ED Now (or PCP triage):**
  - **If No PCP (Primary Care Provider) Second-Level Triage:** You need to be seen within the next hour. Go to the ED/UCC at \_\_\_\_\_ Hospital. Leave as soon as you can.
  - **If PCP Second-Level Triage Required:** You may need to be seen. Your doctor (or NP/PA) will want to talk with you to decide what's best. I'll page the provider on-call now. If you haven't heard from the provider (or me) within 30 minutes, go directly to the ED/UCC at \_\_\_\_\_ Hospital.
- 2 Note to Triager - If NO PCP, Have Other HCP Re-triage the Patient, if Available:**
  - During this COVID-19 pandemic, the medical community is trying to prevent unnecessary referrals to the emergency department (ED). Some patients are fearful of being exposed to COVID-19 in a medical setting. Second-level triage (re-triage) by a physician has been shown to reduce ED referrals. Here are resources that may be available in your community.
  - **PCP Second-level Telephone Triage:** Some PCPs (primary care providers) want to provide re-triage before any of their non-emergent patients are referred to an ED. This requires their approval.
  - **Telemedicine:** Telemedicine is often a preferred source of second-level triage and care during this pandemic. Many practices and some hospitals now offer a telemedicine (virtual visit) service. There are also many national telemedicine companies that are delivering COVID-19 care.
  - **Emergency Department (ED):** Some EDs may provide a telephone follow-up service for patients who have COVID-19 with worsening symptoms.
- 3 Reassurance and Education - Going to the ED or Urgent Care Center During the COVID-19 Pandemic:**
  - If you or your child needs to be seen for an urgent medical problem, do not hesitate to go.
  - ERs and urgent care centers are safe places. They are well equipped to protect you against the virus.
  - For non-urgent conditions, talk to your healthcare provider's office first.
- 4 General Care Advice for COVID-19 Symptoms:**
  - The treatment is the same whether you have COVID-19, influenza or some other respiratory virus.
  - *Cough:* Use cough drops.
  - *Feeling dehydrated:* Drink extra liquids. If the air in your home is dry, use a humidifier.
  - *Fever:* For fever over 101 F (38.3 C), take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package. Do not take aspirin unless your doctor has prescribed it for you.
  - *Muscle aches, headache, and other pains:* Often this comes and goes with the fever. Take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package.
  - *Sore throat:* Try throat lozenges, hard candy or warm chicken broth.
- 5 Care Advice given per Coronavirus (COVID-19) - Diagnosed or Suspected (Adult) guideline.**

## See HCP within 4 Hours (or PCP triage)

Fever > 103 F (39.4 C)

*R/O: serious bacterial infection.*

- 1 See HCP Within 4 Hours (or PCP triage):**
  - **If Office Will Be Open:** You need to be seen within the next 3 or 4 hours. Call your doctor (or NP/PA) now or as soon as the office opens.
  - **If Office Will Be Closed and No PCP (Primary Care Provider) Second-Level Triage:** You need to be seen within the next 3 or 4 hours. A nearby Urgent Care Center (UCC) is often a good source of care. Another choice is to go to the ED. Go sooner if you become worse.
  - **If Office Will Be Closed and PCP Second-Level Triage Required:** You may need to be seen. Your doctor (or NP/PA) will want to talk with you to decide what's best. I'll page the on-call provider now. If you haven't heard from the provider (or me) within 30 minutes, call again. **NOTE:** If on-call provider can't be reached, send to UCC or ED.
- 2 Alternate Disposition - Call Telemedicine Provider Now:**
  - Telemedicine may be your best choice for care during this COVID-19 outbreak.
  - You should call a telemedicine provider now, if your own healthcare provider is not available.

- 3 **Note to Triager - If NO PCP, Have Other HCP Re-triage the Patient, if Available:**
- During this COVID-19 pandemic, the medical community is trying to prevent unnecessary referrals to the emergency department (ED). Some patients are fearful of being exposed to COVID-19 in a medical setting. Second-level triage (re-triage) by a physician has been shown to reduce ED referrals. Here are resources that may be available in your community.
  - **PCP Second-level Telephone Triage:** Some PCPs (primary care providers) want to provide re-triage before any of their non-emergent patients are referred to an ED. This requires their approval.
  - **Telemedicine:** Telemedicine is often a preferred source of second-level triage and care during this pandemic. Many practices and some hospitals now offer a telemedicine (virtual visit) service. There are also many national telemedicine companies that are delivering COVID-19 care.
  - **Emergency Department (ED):** Some EDs may provide a telephone follow-up service for patients who have COVID-19 with worsening symptoms.
- 4 **Note to Triager - Ibuprofen Concerns:**
- Discuss only if caller brings up concerns about ibuprofen.
  - **Response:** The CDC, WHO, and other experts continue to support the use of ibuprofen (if needed) for patients with COVID-19. They found no scientific evidence to support the claim that ibuprofen made COVID-19 worse.
- 5 **How to Protect Others - When You are Sick with COVID-19:**
- **Stay Home a Minimum of 10 Days:** Home isolation is needed for at least 10 days after the symptoms started. Stay home from school or work if you are sick. Do **Not** go to religious services, child care centers, shopping, or other public places. Do **Not** use public transportation (e.g., bus, taxis, ride-sharing). Do **Not** allow any visitors to your home. Leave the house only if you need to seek urgent medical care.
  - **Cover the Cough:** Cough and sneeze into your shirt sleeve or inner elbow. Don't cough into your hand or the air. If available, cough into a tissue and throw it into a trash can.
  - **Wash Hands Often:** Wash hands often with soap and water. After coughing or sneezing are important times. If soap and water are not available, use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry. Avoid touching your eyes, nose, and mouth with unwashed hands.
  - **Wear a Mask:** Wear a facemask when around others. Always wear a facemask (if available) if you have to leave your home (such as going to a medical facility).
  - **Call First if Medical Care Needed:** Call ahead to get approval and careful directions.
- 6 **General Care Advice for COVID-19 Symptoms:**
- The treatment is the same whether you have COVID-19, influenza or some other respiratory virus.
  - **Cough:** Use cough drops.
  - **Feeling dehydrated:** Drink extra liquids. If the air in your home is dry, use a humidifier.
  - **Fever:** For fever over 101 F (38.3 C), take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package. Do not take aspirin unless your doctor has prescribed it for you.
  - **Muscle aches, headache, and other pains:** Often this comes and goes with the fever. Take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package.
  - **Sore throat:** Try throat lozenges, hard candy or warm chicken broth.
- 7 **Cough Medicines:**
- **OTC Cough Syrups:** The most common cough suppressant in OTC cough medications is dextromethorphan. Often the letters "DM" appear in the name.
  - **OTC Cough Drops:** Cough drops can help a lot, especially for mild coughs. They reduce coughing by soothing your irritated throat and removing that tickle sensation in the back of the throat. Cough drops also have the advantage of portability - you can carry them with you.
  - **Home Remedy - Hard Candy:** Hard candy works just as well as medicine-flavored OTC cough drops. People who have diabetes should use sugar-free candy.
  - **Home Remedy - Honey:** This old home remedy has been shown to help decrease coughing at night. The adult dosage is 2 teaspoons (10 ml) at bedtime. Honey should not be given to infants under one year of age.
- 8 **Caution - Dextromethorphan:**
- Do not try to completely suppress coughs that produce mucus and phlegm. Remember that coughing is helpful in bringing up mucus from the lungs and preventing pneumonia.
  - **Research Notes:** Dextromethorphan in some research studies has been shown to reduce the frequency and severity of cough in adults (18 years or older) without significant adverse effects. However, other studies suggest that dextromethorphan is no better than placebo at reducing a cough.
  - **Drug Abuse Potential:** It should be noted that dextromethorphan has become a drug of abuse. This problem is seen most often in adolescents. Overdose symptoms can range from giggling and euphoria to hallucinations and coma.
  - **Contraindicated:** Do not take dextromethorphan if you are taking a monoamine oxidase (MAO) inhibitor now or in the past 2 weeks. Examples of MAO inhibitors include isocarboxazid (Marplan), phenelzine (Nardil), selegiline (Eldepryl, Emsam, Zelapar), and tranylcypromine (Parnate). Do not take dextromethorphan if you are taking venlafaxine (Effexor).



## 9 Fever Medicines:

- For fevers above 101° F (38.3° C) take either acetaminophen or ibuprofen.
- They are over-the-counter (OTC) drugs that help treat both fever and pain. You can buy them at the drugstore.
- The goal of fever therapy is to bring the fever down to a comfortable level. Remember that fever medicine usually lowers fever 2 degrees F (1 - 1 1/2 degrees C).
- **Acetaminophen Regular Strength Tylenol:** Take 650 mg (two 325 mg pills) by mouth every 4-6 hours as needed. Each Regular Strength Tylenol pill has 325 mg of acetaminophen. The most you should take each day is 3,250 mg (10 pills a day).
- **Acetaminophen - Extra Strength Tylenol:** Take 1,000 mg (two 500 mg pills) every 8 hours as needed. Each Extra Strength Tylenol pill has 500 mg of acetaminophen. The most you should take each day is 3,000 mg (6 pills a day).
- **Ibuprofen (e.g., Motrin, Advil):** Take 400 mg (two 200 mg pills) by mouth every 6 hours. The most you should take each day is 1,200 mg (six 200 mg pills), unless your doctor has told you to take more.

## 10 Fever Medicines - Extra Notes and Warnings:

- Use the lowest amount of medicine that makes your fever better.
- Acetaminophen is thought to be safer than ibuprofen or naproxen in people over 65 years old. Acetaminophen is in many OTC and prescription medicines. It might be in more than one medicine that you are taking. You need to be careful and not take an overdose. An acetaminophen overdose can hurt the liver.
- McNeil, the company that makes Tylenol, has different dosage instructions for Tylenol in Canada and the United States. In Canada, the maximum recommended dose per day is 4,000 mg or twelve Regular-Strength (325 mg) pills. In the United States, McNeil recommends a maximum dose of ten Regular-Strength (325 mg) pills.
- **Caution:** Do not take acetaminophen if you have liver disease.
- **Caution:** Do not take ibuprofen if you have stomach problems, kidney disease, are pregnant, or have been told by your doctor to avoid this type of anti-inflammatory drug. Do not take ibuprofen for more than 7 days without consulting your doctor.
- *Before taking any medicine, read all the instructions on the package.*

## 11 Call Back If:

- You become worse.

## 12 Care Advice given per Coronavirus (COVID-19) - Diagnosed or Suspected (Adult) guideline.

[1] Fever > 101 F (38.3 C) AND [2] age > 60

R/O: pneumonia

### 1 See HCP Within 4 Hours (or PCP triage):

- **If Office Will Be Open:** You need to be seen within the next 3 or 4 hours. Call your doctor (or NP/PA) now or as soon as the office opens.
- **If Office Will Be Closed and No PCP (Primary Care Provider) Second-Level Triage:** You need to be seen within the next 3 or 4 hours. A nearby Urgent Care Center (UCC) is often a good source of care. Another choice is to go to the ED. Go sooner if you become worse.
- **If Office Will Be Closed and PCP Second-Level Triage Required:** You may need to be seen. Your doctor (or NP/PA) will want to talk with you to decide what's best. I'll page the on-call provider now. If you haven't heard from the provider (or me) within 30 minutes, call again. **NOTE:** If on-call provider can't be reached, send to UCC or ED.

### 2 Alternate Disposition - Call Telemedicine Provider Now:

- Telemedicine may be your best choice for care during this COVID-19 outbreak.
- You should call a telemedicine provider now, if your own healthcare provider is not available.

### 3 Note to Triager - If NO PCP, Have Other HCP Re-triage the Patient, if Available:

- During this COVID-19 pandemic, the medical community is trying to prevent unnecessary referrals to the emergency department (ED). Some patients are fearful of being exposed to COVID-19 in a medical setting. Second-level triage (re-triage) by a physician has been shown to reduce ED referrals. Here are resources that may be available in your community.
- **PCP Second-level Telephone Triage:** Some PCPs (primary care providers) want to provide re-triage before any of their non-emergent patients are referred to an ED. This requires their approval.
- **Telemedicine:** Telemedicine is often a preferred source of second-level triage and care during this pandemic. Many practices and some hospitals now offer a telemedicine (virtual visit) service. There are also many national telemedicine companies that are delivering COVID-19 care.
- **Emergency Department (ED):** Some EDs may provide a telephone follow-up service for patients who have COVID-19 with worsening symptoms.

### 4 Note to Triager - Ibuprofen Concerns:

- Discuss only if caller brings up concerns about ibuprofen.
- **Response:** The CDC, WHO, and other experts continue to support the use of ibuprofen (if needed) for patients with COVID-19. They found no scientific evidence to support the claim that ibuprofen made COVID-19 worse.

## 5 How to Protect Others - When You are Sick with COVID-19:

- **Stay Home a Minimum of 10 Days:** Home isolation is needed for at least 10 days after the symptoms started. Stay home from school or work if you are sick. Do **Not** go to religious services, child care centers, shopping, or other public places. Do **Not** use public transportation (e.g., bus, taxis, ride-sharing). Do **Not** allow any visitors to your home. Leave the house only if you need to seek urgent medical care.
- **Cover the Cough:** Cough and sneeze into your shirt sleeve or inner elbow. Don't cough into your hand or the air. If available, cough into a tissue and throw it into a trash can.
- **Wash Hands Often:** Wash hands often with soap and water. After coughing or sneezing are important times. If soap and water are not available, use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry. Avoid touching your eyes, nose, and mouth with unwashed hands.
- **Wear a Mask:** Wear a facemask when around others. Always wear a facemask (if available) if you have to leave your home (such as going to a medical facility).
- **Call First if Medical Care Needed:** Call ahead to get approval and careful directions.

## 6 General Care Advice for COVID-19 Symptoms:

- The treatment is the same whether you have COVID-19, influenza or some other respiratory virus.
- **Cough:** Use cough drops.
- **Feeling dehydrated:** Drink extra liquids. If the air in your home is dry, use a humidifier.
- **Fever:** For fever over 101 F (38.3 C), take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package. Do not take aspirin unless your doctor has prescribed it for you.
- **Muscle aches, headache, and other pains:** Often this comes and goes with the fever. Take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package.
- **Sore throat:** Try throat lozenges, hard candy or warm chicken broth.

## 7 Cough Medicines:

- **OTC Cough Syrups:** The most common cough suppressant in OTC cough medications is dextromethorphan. Often the letters "DM" appear in the name.
- **OTC Cough Drops:** Cough drops can help a lot, especially for mild coughs. They reduce coughing by soothing your irritated throat and removing that tickle sensation in the back of the throat. Cough drops also have the advantage of portability - you can carry them with you.
- **Home Remedy - Hard Candy:** Hard candy works just as well as medicine-flavored OTC cough drops. People who have diabetes should use sugar-free candy.
- **Home Remedy - Honey:** This old home remedy has been shown to help decrease coughing at night. The adult dosage is 2 teaspoons (10 ml) at bedtime. Honey should not be given to infants under one year of age.

## 8 Caution - Dextromethorphan:

- Do not try to completely suppress coughs that produce mucus and phlegm. Remember that coughing is helpful in bringing up mucus from the lungs and preventing pneumonia.
- **Research Notes:** Dextromethorphan in some research studies has been shown to reduce the frequency and severity of cough in adults (18 years or older) without significant adverse effects. However, other studies suggest that dextromethorphan is no better than placebo at reducing a cough.
- **Drug Abuse Potential:** It should be noted that dextromethorphan has become a drug of abuse. This problem is seen most often in adolescents. Overdose symptoms can range from giggling and euphoria to hallucinations and coma.
- **Contraindicated:** Do not take dextromethorphan if you are taking a monoamine oxidase (MAO) inhibitor now or in the past 2 weeks. Examples of MAO inhibitors include isocarboxazid (Marplan), phenelzine (Nardil), selegiline (Eldepryl, Emsam, Zelapar), and tranylcypromine (Parnate). Do not take dextromethorphan if you are taking venlafaxine (Effexor).

## 9 Fever Medicines:

- For fevers above 101° F (38.3° C) take either acetaminophen or ibuprofen.
- They are over-the-counter (OTC) drugs that help treat both fever and pain. You can buy them at the drugstore.
- The goal of fever therapy is to bring the fever down to a comfortable level. Remember that fever medicine usually lowers fever 2 degrees F (1 - 1 1/2 degrees C).
- **Acetaminophen Regular Strength Tylenol:** Take 650 mg (two 325 mg pills) by mouth every 4-6 hours as needed. Each Regular Strength Tylenol pill has 325 mg of acetaminophen. The most you should take each day is 3,250 mg (10 pills a day).
- **Acetaminophen - Extra Strength Tylenol:** Take 1,000 mg (two 500 mg pills) every 8 hours as needed. Each Extra Strength Tylenol pill has 500 mg of acetaminophen. The most you should take each day is 3,000 mg (6 pills a day).
- **Ibuprofen (e.g., Motrin, Advil):** Take 400 mg (two 200 mg pills) by mouth every 6 hours. The most you should take each day is 1,200 mg (six 200 mg pills), unless your doctor has told you to take more.

10 **Fever Medicines - Extra Notes and Warnings:**

- Use the lowest amount of medicine that makes your fever better.
- Acetaminophen is thought to be safer than ibuprofen or naproxen in people over 65 years old. Acetaminophen is in many OTC and prescription medicines. It might be in more than one medicine that you are taking. You need to be careful and not take an overdose. An acetaminophen overdose can hurt the liver.
- McNeil, the company that makes Tylenol, has different dosage instructions for Tylenol in Canada and the United States. In Canada, the maximum recommended dose per day is 4,000 mg or twelve Regular-Strength (325 mg) pills. In the United States, McNeil recommends a maximum dose of ten Regular-Strength (325 mg) pills.
- **Caution:** Do not take acetaminophen if you have liver disease.
- **Caution:** Do not take ibuprofen if you have stomach problems, kidney disease, are pregnant, or have been told by your doctor to avoid this type of anti-inflammatory drug. Do not take ibuprofen for more than 7 days without consulting your doctor.
- *Before taking any medicine, read all the instructions on the package.*

11 **Call Back If:**

- You become worse.

12 **Care Advice** given per **Coronavirus (COVID-19) - Diagnosed or Suspected** (Adult) guideline.

[1] Fever > 100.0 F (37.8 C) AND [2] bedridden (e.g., nursing home patient, CVA, chronic illness, recovering from surgery)

*R/O: pneumonia*

1 **See HCP Within 4 Hours (or PCP triage):**

- **If Office Will Be Open:** You need to be seen within the next 3 or 4 hours. Call your doctor (or NP/PA) now or as soon as the office opens.
- **If Office Will Be Closed and No PCP (Primary Care Provider) Second-Level Triage:** You need to be seen within the next 3 or 4 hours. A nearby Urgent Care Center (UCC) is often a good source of care. Another choice is to go to the ED. Go sooner if you become worse.
- **If Office Will Be Closed and PCP Second-Level Triage Required:** You may need to be seen. Your doctor (or NP/PA) will want to talk with you to decide what's best. I'll page the on-call provider now. If you haven't heard from the provider (or me) within 30 minutes, call again. **NOTE:** If on-call provider can't be reached, send to UCC or ED.

2 **Alternate Disposition - Call Telemedicine Provider Now:**

- Telemedicine may be your best choice for care during this COVID-19 outbreak.
- You should call a telemedicine provider now, if your own healthcare provider is not available.

3 **Note to Triager - If NO PCP, Have Other HCP Re-triage the Patient, if Available:**

- During this COVID-19 pandemic, the medical community is trying to prevent unnecessary referrals to the emergency department (ED). Some patients are fearful of being exposed to COVID-19 in a medical setting. Second-level triage (re-triage) by a physician has been shown to reduce ED referrals. Here are resources that may be available in your community.
- **PCP Second-level Telephone Triage:** Some PCPs (primary care providers) want to provide re-triage before any of their non-emergent patients are referred to an ED. This requires their approval.
- **Telemedicine:** Telemedicine is often a preferred source of second-level triage and care during this pandemic. Many practices and some hospitals now offer a telemedicine (virtual visit) service. There are also many national telemedicine companies that are delivering COVID-19 care.
- **Emergency Department (ED):** Some EDs may provide a telephone follow-up service for patients who have COVID-19 with worsening symptoms.

4 **Note to Triager - Ibuprofen Concerns:**

- Discuss only if caller brings up concerns about ibuprofen.
- **Response:** The CDC, WHO, and other experts continue to support the use of ibuprofen (if needed) for patients with COVID-19. They found no scientific evidence to support the claim that ibuprofen made COVID-19 worse.

## 5 How to Protect Others - When You are Sick with COVID-19:

- **Stay Home a Minimum of 10 Days:** Home isolation is needed for at least 10 days after the symptoms started. Stay home from school or work if you are sick. Do **Not** go to religious services, child care centers, shopping, or other public places. Do **Not** use public transportation (e.g., bus, taxis, ride-sharing). Do **Not** allow any visitors to your home. Leave the house only if you need to seek urgent medical care.
- **Cover the Cough:** Cough and sneeze into your shirt sleeve or inner elbow. Don't cough into your hand or the air. If available, cough into a tissue and throw it into a trash can.
- **Wash Hands Often:** Wash hands often with soap and water. After coughing or sneezing are important times. If soap and water are not available, use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry. Avoid touching your eyes, nose, and mouth with unwashed hands.
- **Wear a Mask:** Wear a facemask when around others. Always wear a facemask (if available) if you have to leave your home (such as going to a medical facility).
- **Call First if Medical Care Needed:** Call ahead to get approval and careful directions.

## 6 General Care Advice for COVID-19 Symptoms:

- The treatment is the same whether you have COVID-19, influenza or some other respiratory virus.
- **Cough:** Use cough drops.
- **Feeling dehydrated:** Drink extra liquids. If the air in your home is dry, use a humidifier.
- **Fever:** For fever over 101 F (38.3 C), take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package. Do not take aspirin unless your doctor has prescribed it for you.
- **Muscle aches, headache, and other pains:** Often this comes and goes with the fever. Take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package.
- **Sore throat:** Try throat lozenges, hard candy or warm chicken broth.

## 7 Cough Medicines:

- **OTC Cough Syrups:** The most common cough suppressant in OTC cough medications is dextromethorphan. Often the letters "DM" appear in the name.
- **OTC Cough Drops:** Cough drops can help a lot, especially for mild coughs. They reduce coughing by soothing your irritated throat and removing that tickle sensation in the back of the throat. Cough drops also have the advantage of portability - you can carry them with you.
- **Home Remedy - Hard Candy:** Hard candy works just as well as medicine-flavored OTC cough drops. People who have diabetes should use sugar-free candy.
- **Home Remedy - Honey:** This old home remedy has been shown to help decrease coughing at night. The adult dosage is 2 teaspoons (10 ml) at bedtime. Honey should not be given to infants under one year of age.

## 8 Caution - Dextromethorphan:

- Do not try to completely suppress coughs that produce mucus and phlegm. Remember that coughing is helpful in bringing up mucus from the lungs and preventing pneumonia.
- **Research Notes:** Dextromethorphan in some research studies has been shown to reduce the frequency and severity of cough in adults (18 years or older) without significant adverse effects. However, other studies suggest that dextromethorphan is no better than placebo at reducing a cough.
- **Drug Abuse Potential:** It should be noted that dextromethorphan has become a drug of abuse. This problem is seen most often in adolescents. Overdose symptoms can range from giggling and euphoria to hallucinations and coma.
- **Contraindicated:** Do not take dextromethorphan if you are taking a monoamine oxidase (MAO) inhibitor now or in the past 2 weeks. Examples of MAO inhibitors include isocarboxazid (Marplan), phenelzine (Nardil), selegiline (Eldepryl, Emsam, Zelapar), and tranylcypromine (Parnate). Do not take dextromethorphan if you are taking venlafaxine (Effexor).

## 9 Fever Medicines:

- For fevers above 101° F (38.3° C) take either acetaminophen or ibuprofen.
- They are over-the-counter (OTC) drugs that help treat both fever and pain. You can buy them at the drugstore.
- The goal of fever therapy is to bring the fever down to a comfortable level. Remember that fever medicine usually lowers fever 2 degrees F (1 - 1 1/2 degrees C).
- **Acetaminophen Regular Strength Tylenol:** Take 650 mg (two 325 mg pills) by mouth every 4-6 hours as needed. Each Regular Strength Tylenol pill has 325 mg of acetaminophen. The most you should take each day is 3,250 mg (10 pills a day).
- **Acetaminophen - Extra Strength Tylenol:** Take 1,000 mg (two 500 mg pills) every 8 hours as needed. Each Extra Strength Tylenol pill has 500 mg of acetaminophen. The most you should take each day is 3,000 mg (6 pills a day).
- **Ibuprofen (e.g., Motrin, Advil):** Take 400 mg (two 200 mg pills) by mouth every 6 hours. The most you should take each day is 1,200 mg (six 200 mg pills), unless your doctor has told you to take more.

10 **Fever Medicines - Extra Notes and Warnings:**

- Use the lowest amount of medicine that makes your fever better.
- Acetaminophen is thought to be safer than ibuprofen or naproxen in people over 65 years old. Acetaminophen is in many OTC and prescription medicines. It might be in more than one medicine that you are taking. You need to be careful and not take an overdose. An acetaminophen overdose can hurt the liver.
- McNeil, the company that makes Tylenol, has different dosage instructions for Tylenol in Canada and the United States. In Canada, the maximum recommended dose per day is 4,000 mg or twelve Regular-Strength (325 mg) pills. In the United States, McNeil recommends a maximum dose of ten Regular-Strength (325 mg) pills.
- **Caution:** Do not take acetaminophen if you have liver disease.
- **Caution:** Do not take ibuprofen if you have stomach problems, kidney disease, are pregnant, or have been told by your doctor to avoid this type of anti-inflammatory drug. Do not take ibuprofen for more than 7 days without consulting your doctor.
- *Before taking any medicine, read all the instructions on the package.*

11 **Call Back If:**

- You become worse.

12 **Care Advice** given per **Coronavirus (COVID-19) - Diagnosed or Suspected (Adult)** guideline.

## Call PCP Now

[1] HIGH RISK patient (e.g., age > 64 years, diabetes, heart or lung disease, weak immune system)  
AND [2] new or worsening symptoms

*Note: HIGH RISK is defined in Background area of this guideline.*

1 **Call PCP Now:**

- You need to discuss this with your doctor (or NP/PA).
- I'll page the on-call provider now. If you haven't heard from the provider (or me) within 30 minutes, call again.

2 **Alternate Disposition - Call Telemedicine Provider Now:**

- Telemedicine may be your best choice for care during this COVID-19 outbreak.
- You should call a telemedicine provider now, if your own healthcare provider is not available.

3 **Note to Triage - If NO PCP, Have Other HCP Re-triage the Patient, if Available:**

- During this COVID-19 pandemic, the medical community is trying to prevent unnecessary referrals to the emergency department (ED). Some patients are fearful of being exposed to COVID-19 in a medical setting. Second-level triage (re-triage) by a physician has been shown to reduce ED referrals. Here are resources that may be available in your community.
- **PCP Second-level Telephone Triage:** Some PCPs (primary care providers) want to provide re-triage before any of their non-emergent patients are referred to an ED. This requires their approval.
- **Telemedicine:** Telemedicine is often a preferred source of second-level triage and care during this pandemic. Many practices and some hospitals now offer a telemedicine (virtual visit) service. There are also many national telemedicine companies that are delivering COVID-19 care.
- **Emergency Department (ED):** Some EDs may provide a telephone follow-up service for patients who have COVID-19 with worsening symptoms.

4 **Note to Triage - Ibuprofen Concerns:**

- Discuss only if caller brings up concerns about ibuprofen.
- **Response:** The CDC, WHO, and other experts continue to support the use of ibuprofen (if needed) for patients with COVID-19. They found no scientific evidence to support the claim that ibuprofen made COVID-19 worse.

5 **How to Protect Others - When You are Sick with COVID-19:**

- **Stay Home a Minimum of 10 Days:** Home isolation is needed for at least 10 days after the symptoms started. Stay home from school or work if you are sick. Do **Not** go to religious services, child care centers, shopping, or other public places. Do **Not** use public transportation (e.g., bus, taxis, ride-sharing). Do **Not** allow any visitors to your home. Leave the house only if you need to seek urgent medical care.
- **Cover the Cough:** Cough and sneeze into your shirt sleeve or inner elbow. Don't cough into your hand or the air. If available, cough into a tissue and throw it into a trash can.
- **Wash Hands Often:** Wash hands often with soap and water. After coughing or sneezing are important times. If soap and water are not available, use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry. Avoid touching your eyes, nose, and mouth with unwashed hands.
- **Wear a Mask:** Wear a facemask when around others. Always wear a facemask (if available) if you have to leave your home (such as going to a medical facility).
- **Call First if Medical Care Needed:** Call ahead to get approval and careful directions.



## 6 General Care Advice for COVID-19 Symptoms:

- The treatment is the same whether you have COVID-19, influenza or some other respiratory virus.
- *Cough*: Use cough drops.
- *Feeling dehydrated*: Drink extra liquids. If the air in your home is dry, use a humidifier.
- *Fever*: For fever over 101 F (38.3 C), take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package. Do not take aspirin unless your doctor has prescribed it for you.
- *Muscle aches, headache, and other pains*: Often this comes and goes with the fever. Take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package.
- *Sore throat*: Try throat lozenges, hard candy or warm chicken broth.

## 7 Cough Medicines:

- **OTC Cough Syrups**: The most common cough suppressant in OTC cough medications is dextromethorphan. Often the letters "DM" appear in the name.
- **OTC Cough Drops**: Cough drops can help a lot, especially for mild coughs. They reduce coughing by soothing your irritated throat and removing that tickle sensation in the back of the throat. Cough drops also have the advantage of portability - you can carry them with you.
- **Home Remedy - Hard Candy**: Hard candy works just as well as medicine-flavored OTC cough drops. People who have diabetes should use sugar-free candy.
- **Home Remedy - Honey**: This old home remedy has been shown to help decrease coughing at night. The adult dosage is 2 teaspoons (10 ml) at bedtime. Honey should not be given to infants under one year of age.

## 8 Caution - Dextromethorphan:

- Do not try to completely suppress coughs that produce mucus and phlegm. Remember that coughing is helpful in bringing up mucus from the lungs and preventing pneumonia.
- **Research Notes**: Dextromethorphan in some research studies has been shown to reduce the frequency and severity of cough in adults (18 years or older) without significant adverse effects. However, other studies suggest that dextromethorphan is no better than placebo at reducing a cough.
- **Drug Abuse Potential**: It should be noted that dextromethorphan has become a drug of abuse. This problem is seen most often in adolescents. Overdose symptoms can range from giggling and euphoria to hallucinations and coma.
- **Contraindicated**: Do not take dextromethorphan if you are taking a monoamine oxidase (MAO) inhibitor now or in the past 2 weeks. Examples of MAO inhibitors include isocarboxazid (Marplan), phenelzine (Nardil), selegiline (Eldepryl, Emsam, Zelapar), and tranylcypromine (Parnate). Do not take dextromethorphan if you are taking venlafaxine (Effexor).

## 9 Fever Medicines:

- For fevers above 101° F (38.3° C) take either acetaminophen or ibuprofen.
- They are over-the-counter (OTC) drugs that help treat both fever and pain. You can buy them at the drugstore.
- The goal of fever therapy is to bring the fever down to a comfortable level. Remember that fever medicine usually lowers fever 2 degrees F (1 - 1 1/2 degrees C).
- **Acetaminophen Regular Strength Tylenol**: Take 650 mg (two 325 mg pills) by mouth every 4-6 hours as needed. Each Regular Strength Tylenol pill has 325 mg of acetaminophen. The most you should take each day is 3,250 mg (10 pills a day).
- **Acetaminophen - Extra Strength Tylenol**: Take 1,000 mg (two 500 mg pills) every 8 hours as needed. Each Extra Strength Tylenol pill has 500 mg of acetaminophen. The most you should take each day is 3,000 mg (6 pills a day).
- **Ibuprofen (e.g., Motrin, Advil)**: Take 400 mg (two 200 mg pills) by mouth every 6 hours. The most you should take each day is 1,200 mg (six 200 mg pills), unless your doctor has told you to take more.

## 10 Fever Medicines - Extra Notes and Warnings:

- Use the lowest amount of medicine that makes your fever better.
- Acetaminophen is thought to be safer than ibuprofen or naproxen in people over 65 years old. Acetaminophen is in many OTC and prescription medicines. It might be in more than one medicine that you are taking. You need to be careful and not take an overdose. An acetaminophen overdose can hurt the liver.
- McNeil, the company that makes Tylenol, has different dosage instructions for Tylenol in Canada and the United States. In Canada, the maximum recommended dose per day is 4,000 mg or twelve Regular-Strength (325 mg) pills. In the United States, McNeil recommends a maximum dose of ten Regular-Strength (325 mg) pills.
- **Caution**: Do not take acetaminophen if you have liver disease.
- **Caution**: Do not take ibuprofen if you have stomach problems, kidney disease, are pregnant, or have been told by your doctor to avoid this type of anti-inflammatory drug. Do not take ibuprofen for more than 7 days without consulting your doctor.
- *Before taking any medicine, read all the instructions on the package.*

## 11 Call Back If:

- You become worse.

## 12 Care Advice given per Coronavirus (COVID-19) - Diagnosed or Suspected (Adult) guideline.



[1] HIGH RISK patient AND [2] influenza is widespread in the community AND [3] ONE OR MORE respiratory symptoms: cough, sore throat, runny or stuffy nose

*Reason: During an influenza outbreak, treatment with antiviral influenza medication should be considered for HIGH RISK patients, especially for symptoms present < 48 hours. PCP may wish to phone in a prescription to the pharmacy. Testing for both COVID-19 and influenza may be needed.*

**1 Call PCP Now:**

- You need to discuss this with your doctor (or NP/PA).
- I'll page the on-call provider now. If you haven't heard from the provider (or me) within 30 minutes, call again.

**2 Alternate Disposition - Call Telemedicine Provider Now:**

- Telemedicine may be your best choice for care during this COVID-19 outbreak.
- You should call a telemedicine provider now, if your own healthcare provider is not available.

**3 Note to Triager - If NO PCP, Have Other HCP Re-triage the Patient, if Available:**

- During this COVID-19 pandemic, the medical community is trying to prevent unnecessary referrals to the emergency department (ED). Some patients are fearful of being exposed to COVID-19 in a medical setting. Second-level triage (re-triage) by a physician has been shown to reduce ED referrals. Here are resources that may be available in your community.
- **PCP Second-level Telephone Triage:** Some PCPs (primary care providers) want to provide re-triage before any of their non-emergent patients are referred to an ED. This requires their approval.
- **Telemedicine:** Telemedicine is often a preferred source of second-level triage and care during this pandemic. Many practices and some hospitals now offer a telemedicine (virtual visit) service. There are also many national telemedicine companies that are delivering COVID-19 care.
- **Emergency Department (ED):** Some EDs may provide a telephone follow-up service for patients who have COVID-19 with worsening symptoms.

**4 Note to Triager - Ibuprofen Concerns:**

- Discuss only if caller brings up concerns about ibuprofen.
- **Response:** The CDC, WHO, and other experts continue to support the use of ibuprofen (if needed) for patients with COVID-19. They found no scientific evidence to support the claim that ibuprofen made COVID-19 worse.

**5 How to Protect Others - When You are Sick with COVID-19:**

- **Stay Home a Minimum of 10 Days:** Home isolation is needed for at least 10 days after the symptoms started. Stay home from school or work if you are sick. Do **Not** go to religious services, child care centers, shopping, or other public places. Do **Not** use public transportation (e.g., bus, taxis, ride-sharing). Do **Not** allow any visitors to your home. Leave the house only if you need to seek urgent medical care.
- **Cover the Cough:** Cough and sneeze into your shirt sleeve or inner elbow. Don't cough into your hand or the air. If available, cough into a tissue and throw it into a trash can.
- **Wash Hands Often:** Wash hands often with soap and water. After coughing or sneezing are important times. If soap and water are not available, use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry. Avoid touching your eyes, nose, and mouth with unwashed hands.
- **Wear a Mask:** Wear a facemask when around others. Always wear a facemask (if available) if you have to leave your home (such as going to a medical facility).
- **Call First if Medical Care Needed:** Call ahead to get approval and careful directions.

**6 General Care Advice for COVID-19 Symptoms:**

- The treatment is the same whether you have COVID-19, influenza or some other respiratory virus.
- **Cough:** Use cough drops.
- **Feeling dehydrated:** Drink extra liquids. If the air in your home is dry, use a humidifier.
- **Fever:** For fever over 101 F (38.3 C), take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package. Do not take aspirin unless your doctor has prescribed it for you.
- **Muscle aches, headache, and other pains:** Often this comes and goes with the fever. Take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package.
- **Sore throat:** Try throat lozenges, hard candy or warm chicken broth.

7 **Cough Medicines:**

- **OTC Cough Syrups:** The most common cough suppressant in OTC cough medications is dextromethorphan. Often the letters "DM" appear in the name.
- **OTC Cough Drops:** Cough drops can help a lot, especially for mild coughs. They reduce coughing by soothing your irritated throat and removing that tickle sensation in the back of the throat. Cough drops also have the advantage of portability - you can carry them with you.
- **Home Remedy - Hard Candy:** Hard candy works just as well as medicine-flavored OTC cough drops. People who have diabetes should use sugar-free candy.
- **Home Remedy - Honey:** This old home remedy has been shown to help decrease coughing at night. The adult dosage is 2 teaspoons (10 ml) at bedtime. Honey should not be given to infants under one year of age.

8 **Caution - Dextromethorphan:**

- Do not try to completely suppress coughs that produce mucus and phlegm. Remember that coughing is helpful in bringing up mucus from the lungs and preventing pneumonia.
- **Research Notes:** Dextromethorphan in some research studies has been shown to reduce the frequency and severity of cough in adults (18 years or older) without significant adverse effects. However, other studies suggest that dextromethorphan is no better than placebo at reducing a cough.
- **Drug Abuse Potential:** It should be noted that dextromethorphan has become a drug of abuse. This problem is seen most often in adolescents. Overdose symptoms can range from giggling and euphoria to hallucinations and coma.
- **Contraindicated:** Do not take dextromethorphan if you are taking a monoamine oxidase (MAO) inhibitor now or in the past 2 weeks. Examples of MAO inhibitors include isocarboxazid (Marplan), phenelzine (Nardil), selegiline (Eldepryl, Emsam, Zelapar), and tranylcypromine (Parnate). Do not take dextromethorphan if you are taking venlafaxine (Effexor).

9 **Fever Medicines:**

- For fevers above 101° F (38.3° C) take either acetaminophen or ibuprofen.
- They are over-the-counter (OTC) drugs that help treat both fever and pain. You can buy them at the drugstore.
- The goal of fever therapy is to bring the fever down to a comfortable level. Remember that fever medicine usually lowers fever 2 degrees F (1 - 1 1/2 degrees C).
- **Acetaminophen Regular Strength Tylenol:** Take 650 mg (two 325 mg pills) by mouth every 4-6 hours as needed. Each Regular Strength Tylenol pill has 325 mg of acetaminophen. The most you should take each day is 3,250 mg (10 pills a day).
- **Acetaminophen - Extra Strength Tylenol:** Take 1,000 mg (two 500 mg pills) every 8 hours as needed. Each Extra Strength Tylenol pill has 500 mg of acetaminophen. The most you should take each day is 3,000 mg (6 pills a day).
- **Ibuprofen (e.g., Motrin, Advil):** Take 400 mg (two 200 mg pills) by mouth every 6 hours. The most you should take each day is 1,200 mg (six 200 mg pills), unless your doctor has told you to take more.

10 **Fever Medicines - Extra Notes and Warnings:**

- Use the lowest amount of medicine that makes your fever better.
- Acetaminophen is thought to be safer than ibuprofen or naproxen in people over 65 years old. Acetaminophen is in many OTC and prescription medicines. It might be in more than one medicine that you are taking. You need to be careful and not take an overdose. An acetaminophen overdose can hurt the liver.
- McNeil, the company that makes Tylenol, has different dosage instructions for Tylenol in Canada and the United States. In Canada, the maximum recommended dose per day is 4,000 mg or twelve Regular-Strength (325 mg) pills. In the United States, McNeil recommends a maximum dose of ten Regular-Strength (325 mg) pills.
- **Caution:** Do not take acetaminophen if you have liver disease.
- **Caution:** Do not take ibuprofen if you have stomach problems, kidney disease, are pregnant, or have been told by your doctor to avoid this type of anti-inflammatory drug. Do not take ibuprofen for more than 7 days without consulting your doctor.
- *Before taking any medicine, read all the instructions on the package.*

11 **No Aspirin:**

- Do not use aspirin for treatment of fever or pain.
- **Reason:** there is an association between influenza and Reyes' Syndrome.

12 **Call Back If:**

- You become worse.

13 **Care Advice** given per **Coronavirus (COVID-19) - Diagnosed or Suspected** (Adult) guideline.

[1] HIGH RISK patient AND [2] influenza exposure within the last 7 days AND [3] ONE OR MORE respiratory symptoms: cough, sore throat, runny or stuffy nose

*Reason: During an influenza outbreak, treatment with antiviral influenza medication should be considered for HIGH RISK patients, especially for symptoms present < 48 hours. PCP may wish to phone in a prescription to the pharmacy. Testing for both COVID-19 and influenza may be needed.*

- 1 **Call PCP Now:**
  - You need to discuss this with your doctor (or NP/PA).
  - I'll page the on-call provider now. If you haven't heard from the provider (or me) within 30 minutes, call again.
- 2 **Alternate Disposition - Call Telemedicine Provider Now:**
  - Telemedicine may be your best choice for care during this COVID-19 outbreak.
  - You should call a telemedicine provider now, if your own healthcare provider is not available.
- 3 **Note to Triager - If NO PCP, Have Other HCP Re-triage the Patient, if Available:**
  - During this COVID-19 pandemic, the medical community is trying to prevent unnecessary referrals to the emergency department (ED). Some patients are fearful of being exposed to COVID-19 in a medical setting. Second-level triage (re-triage) by a physician has been shown to reduce ED referrals. Here are resources that may be available in your community.
    - **PCP Second-level Telephone Triage:** Some PCPs (primary care providers) want to provide re-triage before any of their non-emergent patients are referred to an ED. This requires their approval.
    - **Telemedicine:** Telemedicine is often a preferred source of second-level triage and care during this pandemic. Many practices and some hospitals now offer a telemedicine (virtual visit) service. There are also many national telemedicine companies that are delivering COVID-19 care.
    - **Emergency Department (ED):** Some EDs may provide a telephone follow-up service for patients who have COVID-19 with worsening symptoms.
- 4 **Note to Triager - Ibuprofen Concerns:**
  - Discuss only if caller brings up concerns about ibuprofen.
  - **Response:** The CDC, WHO, and other experts continue to support the use of ibuprofen (if needed) for patients with COVID-19. They found no scientific evidence to support the claim that ibuprofen made COVID-19 worse.
- 5 **How to Protect Others - When You are Sick with COVID-19:**
  - **Stay Home a Minimum of 10 Days:** Home isolation is needed for at least 10 days after the symptoms started. Stay home from school or work if you are sick. Do **Not** go to religious services, child care centers, shopping, or other public places. Do **Not** use public transportation (e.g., bus, taxis, ride-sharing). Do **Not** allow any visitors to your home. Leave the house only if you need to seek urgent medical care.
  - **Cover the Cough:** Cough and sneeze into your shirt sleeve or inner elbow. Don't cough into your hand or the air. If available, cough into a tissue and throw it into a trash can.
  - **Wash Hands Often:** Wash hands often with soap and water. After coughing or sneezing are important times. If soap and water are not available, use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry. Avoid touching your eyes, nose, and mouth with unwashed hands.
  - **Wear a Mask:** Wear a facemask when around others. Always wear a facemask (if available) if you have to leave your home (such as going to a medical facility).
  - **Call First if Medical Care Needed:** Call ahead to get approval and careful directions.
- 6 **General Care Advice for COVID-19 Symptoms:**
  - The treatment is the same whether you have COVID-19, influenza or some other respiratory virus.
  - *Cough:* Use cough drops.
  - *Feeling dehydrated:* Drink extra liquids. If the air in your home is dry, use a humidifier.
  - *Fever:* For fever over 101 F (38.3 C), take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package. Do not take aspirin unless your doctor has prescribed it for you.
  - *Muscle aches, headache, and other pains:* Often this comes and goes with the fever. Take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package.
  - *Sore throat:* Try throat lozenges, hard candy or warm chicken broth.
- 7 **Cough Medicines:**
  - **OTC Cough Syrups:** The most common cough suppressant in OTC cough medications is dextromethorphan. Often the letters "DM" appear in the name.
  - **OTC Cough Drops:** Cough drops can help a lot, especially for mild coughs. They reduce coughing by soothing your irritated throat and removing that tickle sensation in the back of the throat. Cough drops also have the advantage of portability - you can carry them with you.
  - **Home Remedy - Hard Candy:** Hard candy works just as well as medicine-flavored OTC cough drops. People who have diabetes should use sugar-free candy.
  - **Home Remedy - Honey:** This old home remedy has been shown to help decrease coughing at night. The adult dosage is 2 teaspoons (10 ml) at bedtime. Honey should not be given to infants under one year of age.

**8 Caution - Dextromethorphan:**

- Do not try to completely suppress coughs that produce mucus and phlegm. Remember that coughing is helpful in bringing up mucus from the lungs and preventing pneumonia.
- Research Notes: Dextromethorphan in some research studies has been shown to reduce the frequency and severity of cough in adults (18 years or older) without significant adverse effects. However, other studies suggest that dextromethorphan is no better than placebo at reducing a cough.
- Drug Abuse Potential: It should be noted that dextromethorphan has become a drug of abuse. This problem is seen most often in adolescents. Overdose symptoms can range from giggling and euphoria to hallucinations and coma.
- **Contraindicated:** Do not take dextromethorphan if you are taking a monoamine oxidase (MAO) inhibitor now or in the past 2 weeks. Examples of MAO inhibitors include isocarboxazid (Marplan), phenelzine (Nardil), selegiline (Eldepryl), Emsam, Zelapar), and tranylcypromine (Parnate). Do not take dextromethorphan if you are taking venlafaxine (Effexor).

**9 Fever Medicines:**

- For fevers above 101° F (38.3° C) take either acetaminophen or ibuprofen.
- They are over-the-counter (OTC) drugs that help treat both fever and pain. You can buy them at the drugstore.
- The goal of fever therapy is to bring the fever down to a comfortable level. Remember that fever medicine usually lowers fever 2 degrees F (1 - 1 1/2 degrees C).
- **Acetaminophen Regular Strength Tylenol:** Take 650 mg (two 325 mg pills) by mouth every 4-6 hours as needed. Each Regular Strength Tylenol pill has 325 mg of acetaminophen. The most you should take each day is 3,250 mg (10 pills a day).
- **Acetaminophen - Extra Strength Tylenol:** Take 1,000 mg (two 500 mg pills) every 8 hours as needed. Each Extra Strength Tylenol pill has 500 mg of acetaminophen. The most you should take each day is 3,000 mg (6 pills a day).
- **Ibuprofen (e.g., Motrin, Advil):** Take 400 mg (two 200 mg pills) by mouth every 6 hours. The most you should take each day is 1,200 mg (six 200 mg pills), unless your doctor has told you to take more.

**10 Fever Medicines - Extra Notes and Warnings:**

- Use the lowest amount of medicine that makes your fever better.
- Acetaminophen is thought to be safer than ibuprofen or naproxen in people over 65 years old. Acetaminophen is in many OTC and prescription medicines. It might be in more than one medicine that you are taking. You need to be careful and not take an overdose. An acetaminophen overdose can hurt the liver.
- McNeil, the company that makes Tylenol, has different dosage instructions for Tylenol in Canada and the United States. In Canada, the maximum recommended dose per day is 4,000 mg or twelve Regular-Strength (325 mg) pills. In the United States, McNeil recommends a maximum dose of ten Regular-Strength (325 mg) pills.
- **Caution:** Do not take acetaminophen if you have liver disease.
- **Caution:** Do not take ibuprofen if you have stomach problems, kidney disease, are pregnant, or have been told by your doctor to avoid this type of anti-inflammatory drug. Do not take ibuprofen for more than 7 days without consulting your doctor.
- *Before taking any medicine, read all the instructions on the package.*

**11 No Aspirin:**

- Do not use aspirin for treatment of fever or pain.
- *Reason:* there is an association between influenza and Reyes' Syndrome.

**12 Call Back If:**

- You become worse.

**13 Care Advice given per Coronavirus (COVID-19) - Diagnosed or Suspected (Adult) guideline.**

**Call PCP within 24 Hours**

Fever present > 3 days (72 hours)

*R/O: bacterial sinusitis, bronchitis, pneumonia*

**1 Call PCP Within 24 Hours:**

- You need to discuss this with your doctor (or NP/PA) within the next 24 hours.
- **If Office Will Be Open:** Call the office when it opens tomorrow morning.
- **If Office Will Be Closed:** I'll page the on-call provider now. **Exception:** from 9 pm to 9 am. Since this isn't urgent, we'll hold the page until morning.

**2 Alternate Disposition - Telemedicine Within 24 Hours:**

- Telemedicine may be your best choice for care during this COVID-19 outbreak.
- You should call a telemedicine provider within the next 24 hours, if your own healthcare provider is not available.

- 3 Note to Triager - If NO PCP, Have Other HCP Re-triage the Patient, if Available:**
- During this COVID-19 pandemic, the medical community is trying to prevent unnecessary referrals to the emergency department (ED). Some patients are fearful of being exposed to COVID-19 in a medical setting. Second-level triage (re-triage) by a physician has been shown to reduce ED referrals. Here are resources that may be available in your community.
  - **PCP Second-level Telephone Triage:** Some PCPs (primary care providers) want to provide re-triage before any of their non-emergent patients are referred to an ED. This requires their approval.
  - **Telemedicine:** Telemedicine is often a preferred source of second-level triage and care during this pandemic. Many practices and some hospitals now offer a telemedicine (virtual visit) service. There are also many national telemedicine companies that are delivering COVID-19 care.
  - **Emergency Department (ED):** Some EDs may provide a telephone follow-up service for patients who have COVID-19 with worsening symptoms.
- 4 Note to Triager - Ibuprofen Concerns:**
- Discuss only if caller brings up concerns about ibuprofen.
  - **Response:** The CDC, WHO, and other experts continue to support the use of ibuprofen (if needed) for patients with COVID-19. They found no scientific evidence to support the claim that ibuprofen made COVID-19 worse.
- 5 How to Protect Others - When You are Sick with COVID-19:**
- **Stay Home a Minimum of 10 Days:** Home isolation is needed for at least 10 days after the symptoms started. Stay home from school or work if you are sick. Do **Not** go to religious services, child care centers, shopping, or other public places. Do **Not** use public transportation (e.g., bus, taxis, ride-sharing). Do **Not** allow any visitors to your home. Leave the house only if you need to seek urgent medical care.
  - **Cover the Cough:** Cough and sneeze into your shirt sleeve or inner elbow. Don't cough into your hand or the air. If available, cough into a tissue and throw it into a trash can.
  - **Wash Hands Often:** Wash hands often with soap and water. After coughing or sneezing are important times. If soap and water are not available, use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry. Avoid touching your eyes, nose, and mouth with unwashed hands.
  - **Wear a Mask:** Wear a facemask when around others. Always wear a facemask (if available) if you have to leave your home (such as going to a medical facility).
  - **Call First if Medical Care Needed:** Call ahead to get approval and careful directions.
- 6 General Care Advice for COVID-19 Symptoms:**
- The treatment is the same whether you have COVID-19, influenza or some other respiratory virus.
  - **Cough:** Use cough drops.
  - **Feeling dehydrated:** Drink extra liquids. If the air in your home is dry, use a humidifier.
  - **Fever:** For fever over 101 F (38.3 C), take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package. Do not take aspirin unless your doctor has prescribed it for you.
  - **Muscle aches, headache, and other pains:** Often this comes and goes with the fever. Take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package.
  - **Sore throat:** Try throat lozenges, hard candy or warm chicken broth.
- 7 Cough Medicines:**
- **OTC Cough Syrups:** The most common cough suppressant in OTC cough medications is dextromethorphan. Often the letters "DM" appear in the name.
  - **OTC Cough Drops:** Cough drops can help a lot, especially for mild coughs. They reduce coughing by soothing your irritated throat and removing that tickle sensation in the back of the throat. Cough drops also have the advantage of portability - you can carry them with you.
  - **Home Remedy - Hard Candy:** Hard candy works just as well as medicine-flavored OTC cough drops. People who have diabetes should use sugar-free candy.
  - **Home Remedy - Honey:** This old home remedy has been shown to help decrease coughing at night. The adult dosage is 2 teaspoons (10 ml) at bedtime. Honey should not be given to infants under one year of age.
- 8 Caution - Dextromethorphan:**
- Do not try to completely suppress coughs that produce mucus and phlegm. Remember that coughing is helpful in bringing up mucus from the lungs and preventing pneumonia.
  - **Research Notes:** Dextromethorphan in some research studies has been shown to reduce the frequency and severity of cough in adults (18 years or older) without significant adverse effects. However, other studies suggest that dextromethorphan is no better than placebo at reducing a cough.
  - **Drug Abuse Potential:** It should be noted that dextromethorphan has become a drug of abuse. This problem is seen most often in adolescents. Overdose symptoms can range from giggling and euphoria to hallucinations and coma.
  - **Contraindicated:** Do not take dextromethorphan if you are taking a monoamine oxidase (MAO) inhibitor now or in the past 2 weeks. Examples of MAO inhibitors include isocarboxazid (Marplan), phenelzine (Nardil), selegiline (Eldepryl, Emsam, Zelapar), and tranylcypromine (Parnate). Do not take dextromethorphan if you are taking venlafaxine (Effexor).



## 9 Fever Medicines:

- For fevers above 101° F (38.3° C) take either acetaminophen or ibuprofen.
- They are over-the-counter (OTC) drugs that help treat both fever and pain. You can buy them at the drugstore.
- The goal of fever therapy is to bring the fever down to a comfortable level. Remember that fever medicine usually lowers fever 2 degrees F (1 - 1 1/2 degrees C).
- **Acetaminophen Regular Strength Tylenol:** Take 650 mg (two 325 mg pills) by mouth every 4-6 hours as needed. Each Regular Strength Tylenol pill has 325 mg of acetaminophen. The most you should take each day is 3,250 mg (10 pills a day).
- **Acetaminophen - Extra Strength Tylenol:** Take 1,000 mg (two 500 mg pills) every 8 hours as needed. Each Extra Strength Tylenol pill has 500 mg of acetaminophen. The most you should take each day is 3,000 mg (6 pills a day).
- **Ibuprofen (e.g., Motrin, Advil):** Take 400 mg (two 200 mg pills) by mouth every 6 hours. The most you should take each day is 1,200 mg (six 200 mg pills), unless your doctor has told you to take more.

## 10 Fever Medicines - Extra Notes and Warnings:

- Use the lowest amount of medicine that makes your fever better.
- Acetaminophen is thought to be safer than ibuprofen or naproxen in people over 65 years old. Acetaminophen is in many OTC and prescription medicines. It might be in more than one medicine that you are taking. You need to be careful and not take an overdose. An acetaminophen overdose can hurt the liver.
- McNeil, the company that makes Tylenol, has different dosage instructions for Tylenol in Canada and the United States. In Canada, the maximum recommended dose per day is 4,000 mg or twelve Regular-Strength (325 mg) pills. In the United States, McNeil recommends a maximum dose of ten Regular-Strength (325 mg) pills.
- **Caution:** Do not take acetaminophen if you have liver disease.
- **Caution:** Do not take ibuprofen if you have stomach problems, kidney disease, are pregnant, or have been told by your doctor to avoid this type of anti-inflammatory drug. Do not take ibuprofen for more than 7 days without consulting your doctor.
- *Before taking any medicine, read all the instructions on the package.*

## 11 Call Back If:

- Fever over 103 F (39.4 C)
- Chest pain or difficulty breathing occurs
- You become worse.

## 12 Care Advice given per Coronavirus (COVID-19) - Diagnosed or Suspected (Adult) guideline.

[1] Fever returns after gone for over 24 hours AND [2] symptoms worse or not improved

*R/O: bacterial sinusitis, bronchitis, pneumonia*

### 1 Call PCP Within 24 Hours:

- You need to discuss this with your doctor (or NP/PA) within the next 24 hours.
- **If Office Will Be Open:** Call the office when it opens tomorrow morning.
- **If Office Will Be Closed:** I'll page the on-call provider now. **Exception:** from 9 pm to 9 am. Since this isn't urgent, we'll hold the page until morning.

### 2 Alternate Disposition - Telemedicine Within 24 Hours:

- Telemedicine may be your best choice for care during this COVID-19 outbreak.
- You should call a telemedicine provider within the next 24 hours, if your own healthcare provider is not available.

### 3 Note to Triager - If NO PCP, Have Other HCP Re-triage the Patient, if Available:

- During this COVID-19 pandemic, the medical community is trying to prevent unnecessary referrals to the emergency department (ED). Some patients are fearful of being exposed to COVID-19 in a medical setting. Second-level triage (re-triage) by a physician has been shown to reduce ED referrals. Here are resources that may be available in your community.
- **PCP Second-level Telephone Triage:** Some PCPs (primary care providers) want to provide re-triage before any of their non-emergent patients are referred to an ED. This requires their approval.
- **Telemedicine:** Telemedicine is often a preferred source of second-level triage and care during this pandemic. Many practices and some hospitals now offer a telemedicine (virtual visit) service. There are also many national telemedicine companies that are delivering COVID-19 care.
- **Emergency Department (ED):** Some EDs may provide a telephone follow-up service for patients who have COVID-19 with worsening symptoms.

### 4 Note to Triager - Ibuprofen Concerns:

- Discuss only if caller brings up concerns about ibuprofen.
- **Response:** The CDC, WHO, and other experts continue to support the use of ibuprofen (if needed) for patients with COVID-19. They found no scientific evidence to support the claim that ibuprofen made COVID-19 worse.



5 **How to Protect Others - When You are Sick with COVID-19:**

- **Stay Home a Minimum of 10 Days:** Home isolation is needed for at least 10 days after the symptoms started. Stay home from school or work if you are sick. Do **Not** go to religious services, child care centers, shopping, or other public places. Do **Not** use public transportation (e.g., bus, taxis, ride-sharing). Do **Not** allow any visitors to your home. Leave the house only if you need to seek urgent medical care.
- **Cover the Cough:** Cough and sneeze into your shirt sleeve or inner elbow. Don't cough into your hand or the air. If available, cough into a tissue and throw it into a trash can.
- **Wash Hands Often:** Wash hands often with soap and water. After coughing or sneezing are important times. If soap and water are not available, use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry. Avoid touching your eyes, nose, and mouth with unwashed hands.
- **Wear a Mask:** Wear a facemask when around others. Always wear a facemask (if available) if you have to leave your home (such as going to a medical facility).
- **Call First if Medical Care Needed:** Call ahead to get approval and careful directions.

6 **General Care Advice for COVID-19 Symptoms:**

- The treatment is the same whether you have COVID-19, influenza or some other respiratory virus.
- **Cough:** Use cough drops.
- **Feeling dehydrated:** Drink extra liquids. If the air in your home is dry, use a humidifier.
- **Fever:** For fever over 101 F (38.3 C), take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package. Do not take aspirin unless your doctor has prescribed it for you.
- **Muscle aches, headache, and other pains:** Often this comes and goes with the fever. Take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package.
- **Sore throat:** Try throat lozenges, hard candy or warm chicken broth.

7 **Cough Medicines:**

- **OTC Cough Syrups:** The most common cough suppressant in OTC cough medications is dextromethorphan. Often the letters "DM" appear in the name.
- **OTC Cough Drops:** Cough drops can help a lot, especially for mild coughs. They reduce coughing by soothing your irritated throat and removing that tickle sensation in the back of the throat. Cough drops also have the advantage of portability - you can carry them with you.
- **Home Remedy - Hard Candy:** Hard candy works just as well as medicine-flavored OTC cough drops. People who have diabetes should use sugar-free candy.
- **Home Remedy - Honey:** This old home remedy has been shown to help decrease coughing at night. The adult dosage is 2 teaspoons (10 ml) at bedtime. Honey should not be given to infants under one year of age.

8 **Caution - Dextromethorphan:**

- Do not try to completely suppress coughs that produce mucus and phlegm. Remember that coughing is helpful in bringing up mucus from the lungs and preventing pneumonia.
- **Research Notes:** Dextromethorphan in some research studies has been shown to reduce the frequency and severity of cough in adults (18 years or older) without significant adverse effects. However, other studies suggest that dextromethorphan is no better than placebo at reducing a cough.
- **Drug Abuse Potential:** It should be noted that dextromethorphan has become a drug of abuse. This problem is seen most often in adolescents. Overdose symptoms can range from giggling and euphoria to hallucinations and coma.
- **Contraindicated:** Do not take dextromethorphan if you are taking a monoamine oxidase (MAO) inhibitor now or in the past 2 weeks. Examples of MAO inhibitors include isocarboxazid (Marplan), phenelzine (Nardil), selegiline (Eldepryl, Emsam, Zelapar), and tranylcypromine (Parnate). Do not take dextromethorphan if you are taking venlafaxine (Effexor).

9 **Fever Medicines:**

- For fevers above 101° F (38.3° C) take either acetaminophen or ibuprofen.
- They are over-the-counter (OTC) drugs that help treat both fever and pain. You can buy them at the drugstore.
- The goal of fever therapy is to bring the fever down to a comfortable level. Remember that fever medicine usually lowers fever 2 degrees F (1 - 1 1/2 degrees C).
- **Acetaminophen Regular Strength Tylenol:** Take 650 mg (two 325 mg pills) by mouth every 4-6 hours as needed. Each Regular Strength Tylenol pill has 325 mg of acetaminophen. The most you should take each day is 3,250 mg (10 pills a day).
- **Acetaminophen - Extra Strength Tylenol:** Take 1,000 mg (two 500 mg pills) every 8 hours as needed. Each Extra Strength Tylenol pill has 500 mg of acetaminophen. The most you should take each day is 3,000 mg (6 pills a day).
- **Ibuprofen (e.g., Motrin, Advil):** Take 400 mg (two 200 mg pills) by mouth every 6 hours. The most you should take each day is 1,200 mg (six 200 mg pills), unless your doctor has told you to take more.

- 10 **Fever Medicines - Extra Notes and Warnings:**
  - Use the lowest amount of medicine that makes your fever better.
  - Acetaminophen is thought to be safer than ibuprofen or naproxen in people over 65 years old. Acetaminophen is in many OTC and prescription medicines. It might be in more than one medicine that you are taking. You need to be careful and not take an overdose. An acetaminophen overdose can hurt the liver.
  - McNeil, the company that makes Tylenol, has different dosage instructions for Tylenol in Canada and the United States. In Canada, the maximum recommended dose per day is 4,000 mg or twelve Regular-Strength (325 mg) pills. In the United States, McNeil recommends a maximum dose of ten Regular-Strength (325 mg) pills.
  - **Caution:** Do not take acetaminophen if you have liver disease.
  - **Caution:** Do not take ibuprofen if you have stomach problems, kidney disease, are pregnant, or have been told by your doctor to avoid this type of anti-inflammatory drug. Do not take ibuprofen for more than 7 days without consulting your doctor.
  - *Before taking any medicine, read all the instructions on the package.*
- 11 **Call Back If:**
  - Fever over 103 F (39.4 C)
  - Chest pain or difficulty breathing occurs
  - You become worse.
- 12 **Care Advice given per Coronavirus (COVID-19) - Diagnosed or Suspected (Adult) guideline.**

[1] Continuous (nonstop) coughing interferes with work or school AND [2] no improvement using cough treatment per protocol

- 1 **Call PCP Within 24 Hours:**
  - You need to discuss this with your doctor (or NP/PA) within the next 24 hours.
  - **If Office Will Be Open:** Call the office when it opens tomorrow morning.
  - **If Office Will Be Closed:** I'll page the on-call provider now. **Exception:** from 9 pm to 9 am. Since this isn't urgent, we'll hold the page until morning.
- 2 **Alternate Disposition - Telemedicine Within 24 Hours:**
  - Telemedicine may be your best choice for care during this COVID-19 outbreak.
  - You should call a telemedicine provider within the next 24 hours, if your own healthcare provider is not available.
- 3 **Note to Triage - If NO PCP, Have Other HCP Re-triage the Patient, if Available:**
  - During this COVID-19 pandemic, the medical community is trying to prevent unnecessary referrals to the emergency department (ED). Some patients are fearful of being exposed to COVID-19 in a medical setting. Second-level triage (re-triage) by a physician has been shown to reduce ED referrals. Here are resources that may be available in your community.
  - **PCP Second-level Telephone Triage:** Some PCPs (primary care providers) want to provide re-triage before any of their non-emergent patients are referred to an ED. This requires their approval.
  - **Telemedicine:** Telemedicine is often a preferred source of second-level triage and care during this pandemic. Many practices and some hospitals now offer a telemedicine (virtual visit) service. There are also many national telemedicine companies that are delivering COVID-19 care.
  - **Emergency Department (ED):** Some EDs may provide a telephone follow-up service for patients who have COVID-19 with worsening symptoms.
- 4 **Note to Triage - Ibuprofen Concerns:**
  - Discuss only if caller brings up concerns about ibuprofen.
  - **Response:** The CDC, WHO, and other experts continue to support the use of ibuprofen (if needed) for patients with COVID-19. They found no scientific evidence to support the claim that ibuprofen made COVID-19 worse.
- 5 **How to Protect Others - When You are Sick with COVID-19:**
  - **Stay Home a Minimum of 10 Days:** Home isolation is needed for at least 10 days after the symptoms started. Stay home from school or work if you are sick. Do **Not** go to religious services, child care centers, shopping, or other public places. Do **Not** use public transportation (e.g., bus, taxis, ride-sharing). Do **Not** allow any visitors to your home. Leave the house only if you need to seek urgent medical care.
  - **Cover the Cough:** Cough and sneeze into your shirt sleeve or inner elbow. Don't cough into your hand or the air. If available, cough into a tissue and throw it into a trash can.
  - **Wash Hands Often:** Wash hands often with soap and water. After coughing or sneezing are important times. If soap and water are not available, use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry. Avoid touching your eyes, nose, and mouth with unwashed hands.
  - **Wear a Mask:** Wear a facemask when around others. Always wear a facemask (if available) if you have to leave your home (such as going to a medical facility).
  - **Call First if Medical Care Needed:** Call ahead to get approval and careful directions.

## 6 General Care Advice for COVID-19 Symptoms:

- The treatment is the same whether you have COVID-19, influenza or some other respiratory virus.
- *Cough*: Use cough drops.
- *Feeling dehydrated*: Drink extra liquids. If the air in your home is dry, use a humidifier.
- *Fever*: For fever over 101 F (38.3 C), take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package. Do not take aspirin unless your doctor has prescribed it for you.
- *Muscle aches, headache, and other pains*: Often this comes and goes with the fever. Take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package.
- *Sore throat*: Try throat lozenges, hard candy or warm chicken broth.

## 7 Cough Medicines:

- **OTC Cough Syrups**: The most common cough suppressant in OTC cough medications is dextromethorphan. Often the letters "DM" appear in the name.
- **OTC Cough Drops**: Cough drops can help a lot, especially for mild coughs. They reduce coughing by soothing your irritated throat and removing that tickle sensation in the back of the throat. Cough drops also have the advantage of portability - you can carry them with you.
- **Home Remedy - Hard Candy**: Hard candy works just as well as medicine-flavored OTC cough drops. People who have diabetes should use sugar-free candy.
- **Home Remedy - Honey**: This old home remedy has been shown to help decrease coughing at night. The adult dosage is 2 teaspoons (10 ml) at bedtime. Honey should not be given to infants under one year of age.

## 8 Caution - Dextromethorphan:

- Do not try to completely suppress coughs that produce mucus and phlegm. Remember that coughing is helpful in bringing up mucus from the lungs and preventing pneumonia.
- **Research Notes**: Dextromethorphan in some research studies has been shown to reduce the frequency and severity of cough in adults (18 years or older) without significant adverse effects. However, other studies suggest that dextromethorphan is no better than placebo at reducing a cough.
- **Drug Abuse Potential**: It should be noted that dextromethorphan has become a drug of abuse. This problem is seen most often in adolescents. Overdose symptoms can range from giggling and euphoria to hallucinations and coma.
- **Contraindicated**: Do not take dextromethorphan if you are taking a monoamine oxidase (MAO) inhibitor now or in the past 2 weeks. Examples of MAO inhibitors include isocarboxazid (Marplan), phenelzine (Nardil), selegiline (Eldepryl, Emsam, Zelapar), and tranylcypromine (Parnate). Do not take dextromethorphan if you are taking venlafaxine (Effexor).

## 9 Fever Medicines:

- For fevers above 101° F (38.3° C) take either acetaminophen or ibuprofen.
- They are over-the-counter (OTC) drugs that help treat both fever and pain. You can buy them at the drugstore.
- The goal of fever therapy is to bring the fever down to a comfortable level. Remember that fever medicine usually lowers fever 2 degrees F (1 - 1 1/2 degrees C).
- **Acetaminophen Regular Strength Tylenol**: Take 650 mg (two 325 mg pills) by mouth every 4-6 hours as needed. Each Regular Strength Tylenol pill has 325 mg of acetaminophen. The most you should take each day is 3,250 mg (10 pills a day).
- **Acetaminophen - Extra Strength Tylenol**: Take 1,000 mg (two 500 mg pills) every 8 hours as needed. Each Extra Strength Tylenol pill has 500 mg of acetaminophen. The most you should take each day is 3,000 mg (6 pills a day).
- **Ibuprofen (e.g., Motrin, Advil)**: Take 400 mg (two 200 mg pills) by mouth every 6 hours. The most you should take each day is 1,200 mg (six 200 mg pills), unless your doctor has told you to take more.

## 10 Fever Medicines - Extra Notes and Warnings:

- Use the lowest amount of medicine that makes your fever better.
- Acetaminophen is thought to be safer than ibuprofen or naproxen in people over 65 years old. Acetaminophen is in many OTC and prescription medicines. It might be in more than one medicine that you are taking. You need to be careful and not take an overdose. An acetaminophen overdose can hurt the liver.
- McNeil, the company that makes Tylenol, has different dosage instructions for Tylenol in Canada and the United States. In Canada, the maximum recommended dose per day is 4,000 mg or twelve Regular-Strength (325 mg) pills. In the United States, McNeil recommends a maximum dose of ten Regular-Strength (325 mg) pills.
- **Caution**: Do not take acetaminophen if you have liver disease.
- **Caution**: Do not take ibuprofen if you have stomach problems, kidney disease, are pregnant, or have been told by your doctor to avoid this type of anti-inflammatory drug. Do not take ibuprofen for more than 7 days without consulting your doctor.
- *Before taking any medicine, read all the instructions on the package.*

## 11 Call Back If:

- Fever over 103 F (39.4 C)
- Chest pain or difficulty breathing occurs
- You become worse.

12 **Care Advice** given per **Coronavirus (COVID-19) - Diagnosed or Suspected (Adult)** guideline.

### Call PCP when Office is Open

[1] COVID-19 infection suspected by caller or triager AND [2] mild symptoms (cough, fever, or others) AND [3] no complications or SOB

*Reason: PCP will discuss suspected diagnosis and need for testing*

**1 Call PCP When Office Is Open:**

- You need to discuss this with your doctor (or NP/PA) within the next few days.
- Call the office when it is open.

**2 Reassurance and Education - Suspected COVID-19:**

- You suspect you have COVID-19 because you have symptoms that match and you were either exposed to someone with it or because it is widespread in your community.
- Most people who get COVID-19 will have mild illness, can recover at home without medical care, and may not need to be tested. From what you have told me, your symptoms are mild. That is reassuring.
- Talk with your healthcare provider about your symptoms. Ask if testing is needed.
- Call them during regular office hours.
- *Here's some care advice to help you and to help prevent others from getting sick.*

**3 General Care Advice for COVID-19 Symptoms:**

- The treatment is the same whether you have COVID-19, influenza or some other respiratory virus.
- *Cough:* Use cough drops.
- *Feeling dehydrated:* Drink extra liquids. If the air in your home is dry, use a humidifier.
- *Fever:* For fever over 101 F (38.3 C), take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package. Do not take aspirin unless your doctor has prescribed it for you.
- *Muscle aches, headache, and other pains:* Often this comes and goes with the fever. Take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package.
- *Sore throat:* Try throat lozenges, hard candy or warm chicken broth.

**4 Cough Medicines:**

- **OTC Cough Syrups:** The most common cough suppressant in OTC cough medications is dextromethorphan. Often the letters "DM" appear in the name.
- **OTC Cough Drops:** Cough drops can help a lot, especially for mild coughs. They reduce coughing by soothing your irritated throat and removing that tickle sensation in the back of the throat. Cough drops also have the advantage of portability - you can carry them with you.
- **Home Remedy - Hard Candy:** Hard candy works just as well as medicine-flavored OTC cough drops. People who have diabetes should use sugar-free candy.
- **Home Remedy - Honey:** This old home remedy has been shown to help decrease coughing at night. The adult dosage is 2 teaspoons (10 ml) at bedtime. Honey should not be given to infants under one year of age.

**5 Caution - Dextromethorphan:**

- Do not try to completely suppress coughs that produce mucus and phlegm. Remember that coughing is helpful in bringing up mucus from the lungs and preventing pneumonia.
- **Research Notes:** Dextromethorphan in some research studies has been shown to reduce the frequency and severity of cough in adults (18 years or older) without significant adverse effects. However, other studies suggest that dextromethorphan is no better than placebo at reducing a cough.
- **Drug Abuse Potential:** It should be noted that dextromethorphan has become a drug of abuse. This problem is seen most often in adolescents. Overdose symptoms can range from giggling and euphoria to hallucinations and coma.
- **Contraindicated:** Do not take dextromethorphan if you are taking a monoamine oxidase (MAO) inhibitor now or in the past 2 weeks. Examples of MAO inhibitors include isocarboxazid (Marplan), phenelzine (Nardil), selegiline (Eldepryl, Emsam, Zelapar), and tranylcypromine (Parnate). Do not take dextromethorphan if you are taking venlafaxine (Effexor).

**6 Humidifier:**

- If the air is dry, use a humidifier in the bedroom.
- Dry air makes coughs worse.

**7 Coughing Spells:**

- Drink warm fluids. Inhale warm mist. (Reason: both relax the airway and loosen up the phlegm)
- Suck on cough drops or hard candy to coat the irritated throat.

**8 Avoid Tobacco Smoke:**

- Avoid tobacco smoke.
- Smoking or being exposed to smoke makes coughs much worse.

**9 Pain and Fever Medicines:**

- For pain or fever relief, take either acetaminophen or ibuprofen.
- They are over-the-counter (OTC) drugs that help treat both fever and pain. You can buy them at the drugstore.
- Treat fevers above 101° F (38.3° C). The goal of fever therapy is to bring the fever down to a comfortable level. Remember that fever medicine usually lowers fever 2 degrees F (1 - 1 1/2 degrees C).
- **Acetaminophen Regular Strength Tylenol:** Take 650 mg (two 325 mg pills) by mouth every 4 to 6 hours as needed. Each Regular Strength Tylenol pill has 325 mg of acetaminophen. The most you should take each day is 3,250 mg (10 pills a day).
- **Acetaminophen - Extra Strength Tylenol:** Take 1,000 mg (two 500 mg pills) every 8 hours as needed. Each Extra Strength Tylenol pill has 500 mg of acetaminophen. The most you should take each day is 3,000 mg (6 pills a day).
- **Ibuprofen (e.g., Motrin, Advil):** Take 400 mg (two 200 mg pills) by mouth every 6 hours. The most you should take each day is 1,200 mg (six 200 mg pills), unless your doctor has told you to take more.

**10 Fever Medicines - Extra Notes and Warnings:**

- Use the lowest amount of medicine that makes your fever better.
- Acetaminophen is thought to be safer than ibuprofen or naproxen in people over 65 years old. Acetaminophen is in many OTC and prescription medicines. It might be in more than one medicine that you are taking. You need to be careful and not take an overdose. An acetaminophen overdose can hurt the liver.
- McNeil, the company that makes Tylenol, has different dosage instructions for Tylenol in Canada and the United States. In Canada, the maximum recommended dose per day is 4,000 mg or twelve Regular-Strength (325 mg) pills. In the United States, McNeil recommends a maximum dose of ten Regular-Strength (325 mg) pills.
- **Caution:** Do not take acetaminophen if you have liver disease.
- **Caution:** Do not take ibuprofen if you have stomach problems, kidney disease, are pregnant, or have been told by your doctor to avoid this type of anti-inflammatory drug. Do not take ibuprofen for more than 7 days without consulting your doctor.
- *Before taking any medicine, read all the instructions on the package.*

**11 Note to Triager - Ibuprofen Concerns:**

- Discuss only if caller brings up concerns about ibuprofen.
- **Response:** The CDC, WHO, and other experts continue to support the use of ibuprofen (if needed) for patients with COVID-19. They found no scientific evidence to support the claim that ibuprofen made COVID-19 worse.

**12 How to Protect Others - When You are Sick with COVID-19:**

- **Stay Home a Minimum of 10 Days:** Home isolation is needed for at least 10 days after the symptoms started. Stay home from school or work if you are sick. Do **Not** go to religious services, child care centers, shopping, or other public places. Do **Not** use public transportation (e.g., bus, taxis, ride-sharing). Do **Not** allow any visitors to your home. Leave the house only if you need to seek urgent medical care.
- **Cover the Cough:** Cough and sneeze into your shirt sleeve or inner elbow. Don't cough into your hand or the air. If available, cough into a tissue and throw it into a trash can.
- **Wash Hands Often:** Wash hands often with soap and water. After coughing or sneezing are important times. If soap and water are not available, use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry. Avoid touching your eyes, nose, and mouth with unwashed hands.
- **Wear a Mask:** Wear a facemask when around others. Always wear a facemask (if available) if you have to leave your home (such as going to a medical facility).
- **Call First if Medical Care Needed:** Call ahead to get approval and careful directions.

**13 Stopping Home Isolation - Must Meet all 3 Requirements (CDC):**

- Fever gone for at least 24 hours off fever-reducing medicines **AND**
- Cough and other symptoms must be improved **AND**
- Symptoms started more than 10 days ago.
- *If unsure if it is safe for you to leave isolation, check the CDC website or call your healthcare provider.*

14 **Other COVID-19 Facts:**

- **Incubation Period:** Average 5 days (range 2 to 14 days) after coming in contact with a person who has COVID-19 virus.
- **No Symptoms, but Infected (Asymptomatic):** Approximately 30% of infected patients may have no symptoms.
- **Mild Infections:** About 80% of those with symptoms have a mild illness, much like a normal flu or a bad cold. The symptoms usually last 2 weeks.
- **Severe Infections:** About 20% of those with symptoms develop trouble breathing from viral pneumonia. Many of these need to be admitted to the hospital. People with complications generally recover in 3 to 6 weeks.
- **Death Rate:** The adult death rate is approximately 1% to 3%. The death rate is lower in children and younger adults. It is higher in older adults.
- **Treatment:** Treatment is supportive. Oxygen and IV fluids are used for hospitalized patients. On May 1, 2020 the U.S. Food and Drug Administration (FDA) granted temporary emergency use authorization (EUA) for the investigational antiviral remdesivir to treat COVID-19. Doctors can use this drug to treat patients who are hospitalized with COVID-19 who are severely ill. More research is needed to determine how well this drug works and for which patients.
- **Prevention - Vaccine:** There currently is no approved vaccine to prevent COVID-19. There are vaccines in development. A COVID-19 vaccine may be available for adults by early 2021.
- **Prevention - Medicine:** Currently, there is no medicine to prevent COVID-19. Warning: the malaria drug (chloroquine) was studied and found not to be helpful for this disease. It also had cardiac side effects. *Social distancing and wearing masks have been proven to help prevent COVID-19!*

15 **Call Back If:**

- Fever over 103 F (39.4 C)
- Fever lasts over 3 days
- Fever returns after being gone for 24 hours
- Chest pain or difficulty breathing occurs
- You become worse.

16 **Care Advice** given per **Coronavirus (COVID-19) - Diagnosed or Suspected** (Adult) guideline.

Cough present > 3 weeks

1 **Call PCP When Office Is Open:**

- You need to discuss this with your doctor (or NP/PA) within the next few days.
- Call the office when it is open.

2 **Alternate Disposition - Telemedicine Within 24 Hours:**

- Telemedicine may be your best choice for care during this COVID-19 outbreak.
- You should call a telemedicine provider within the next 24 hours, if your own healthcare provider is not available.

3 **Note to Triage - If NO PCP, Have Other HCP Re-triage the Patient, if Available:**

- During this COVID-19 pandemic, the medical community is trying to prevent unnecessary referrals to the emergency department (ED). Some patients are fearful of being exposed to COVID-19 in a medical setting. Second-level triage (re-triage) by a physician has been shown to reduce ED referrals. Here are resources that may be available in your community.
- **PCP Second-level Telephone Triage:** Some PCPs (primary care providers) want to provide re-triage before any of their non-emergent patients are referred to an ED. This requires their approval.
- **Telemedicine:** Telemedicine is often a preferred source of second-level triage and care during this pandemic. Many practices and some hospitals now offer a telemedicine (virtual visit) service. There are also many national telemedicine companies that are delivering COVID-19 care.
- **Emergency Department (ED):** Some EDs may provide a telephone follow-up service for patients who have COVID-19 with worsening symptoms.

4 **Note to Triage - Ibuprofen Concerns:**

- Discuss only if caller brings up concerns about ibuprofen.
- **Response:** The CDC, WHO, and other experts continue to support the use of ibuprofen (if needed) for patients with COVID-19. They found no scientific evidence to support the claim that ibuprofen made COVID-19 worse.



## 5 How to Protect Others - When You are Sick with COVID-19:

- **Stay Home a Minimum of 10 Days:** Home isolation is needed for at least 10 days after the symptoms started. Stay home from school or work if you are sick. Do **Not** go to religious services, child care centers, shopping, or other public places. Do **Not** use public transportation (e.g., bus, taxis, ride-sharing). Do **Not** allow any visitors to your home. Leave the house only if you need to seek urgent medical care.
- **Cover the Cough:** Cough and sneeze into your shirt sleeve or inner elbow. Don't cough into your hand or the air. If available, cough into a tissue and throw it into a trash can.
- **Wash Hands Often:** Wash hands often with soap and water. After coughing or sneezing are important times. If soap and water are not available, use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry. Avoid touching your eyes, nose, and mouth with unwashed hands.
- **Wear a Mask:** Wear a facemask when around others. Always wear a facemask (if available) if you have to leave your home (such as going to a medical facility).
- **Call First if Medical Care Needed:** Call ahead to get approval and careful directions.

## 6 General Care Advice for COVID-19 Symptoms:

- The treatment is the same whether you have COVID-19, influenza or some other respiratory virus.
- **Cough:** Use cough drops.
- **Feeling dehydrated:** Drink extra liquids. If the air in your home is dry, use a humidifier.
- **Fever:** For fever over 101 F (38.3 C), take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package. Do not take aspirin unless your doctor has prescribed it for you.
- **Muscle aches, headache, and other pains:** Often this comes and goes with the fever. Take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package.
- **Sore throat:** Try throat lozenges, hard candy or warm chicken broth.

## 7 Cough Medicines:

- **OTC Cough Syrups:** The most common cough suppressant in OTC cough medications is dextromethorphan. Often the letters "DM" appear in the name.
- **OTC Cough Drops:** Cough drops can help a lot, especially for mild coughs. They reduce coughing by soothing your irritated throat and removing that tickle sensation in the back of the throat. Cough drops also have the advantage of portability - you can carry them with you.
- **Home Remedy - Hard Candy:** Hard candy works just as well as medicine-flavored OTC cough drops. People who have diabetes should use sugar-free candy.
- **Home Remedy - Honey:** This old home remedy has been shown to help decrease coughing at night. The adult dosage is 2 teaspoons (10 ml) at bedtime. Honey should not be given to infants under one year of age.

## 8 Caution - Dextromethorphan:

- Do not try to completely suppress coughs that produce mucus and phlegm. Remember that coughing is helpful in bringing up mucus from the lungs and preventing pneumonia.
- **Research Notes:** Dextromethorphan in some research studies has been shown to reduce the frequency and severity of cough in adults (18 years or older) without significant adverse effects. However, other studies suggest that dextromethorphan is no better than placebo at reducing a cough.
- **Drug Abuse Potential:** It should be noted that dextromethorphan has become a drug of abuse. This problem is seen most often in adolescents. Overdose symptoms can range from giggling and euphoria to hallucinations and coma.
- **Contraindicated:** Do not take dextromethorphan if you are taking a monoamine oxidase (MAO) inhibitor now or in the past 2 weeks. Examples of MAO inhibitors include isocarboxazid (Marplan), phenelzine (Nardil), selegiline (Eldepryl, Emsam, Zelapar), and tranylcypromine (Parnate). Do not take dextromethorphan if you are taking venlafaxine (Effexor).

## 9 Fever Medicines:

- For fevers above 101° F (38.3° C) take either acetaminophen or ibuprofen.
- They are over-the-counter (OTC) drugs that help treat both fever and pain. You can buy them at the drugstore.
- The goal of fever therapy is to bring the fever down to a comfortable level. Remember that fever medicine usually lowers fever 2 degrees F (1 - 1 1/2 degrees C).
- **Acetaminophen Regular Strength Tylenol:** Take 650 mg (two 325 mg pills) by mouth every 4-6 hours as needed. Each Regular Strength Tylenol pill has 325 mg of acetaminophen. The most you should take each day is 3,250 mg (10 pills a day).
- **Acetaminophen - Extra Strength Tylenol:** Take 1,000 mg (two 500 mg pills) every 8 hours as needed. Each Extra Strength Tylenol pill has 500 mg of acetaminophen. The most you should take each day is 3,000 mg (6 pills a day).
- **Ibuprofen (e.g., Motrin, Advil):** Take 400 mg (two 200 mg pills) by mouth every 6 hours. The most you should take each day is 1,200 mg (six 200 mg pills), unless your doctor has told you to take more.

10 **Fever Medicines - Extra Notes and Warnings:**

- Use the lowest amount of medicine that makes your fever better.
- Acetaminophen is thought to be safer than ibuprofen or naproxen in people over 65 years old. Acetaminophen is in many OTC and prescription medicines. It might be in more than one medicine that you are taking. You need to be careful and not take an overdose. An acetaminophen overdose can hurt the liver.
- McNeil, the company that makes Tylenol, has different dosage instructions for Tylenol in Canada and the United States. In Canada, the maximum recommended dose per day is 4,000 mg or twelve Regular-Strength (325 mg) pills. In the United States, McNeil recommends a maximum dose of ten Regular-Strength (325 mg) pills.
- **Caution:** Do not take acetaminophen if you have liver disease.
- **Caution:** Do not take ibuprofen if you have stomach problems, kidney disease, are pregnant, or have been told by your doctor to avoid this type of anti-inflammatory drug. Do not take ibuprofen for more than 7 days without consulting your doctor.
- *Before taking any medicine, read all the instructions on the package.*

11 **Call Back If:**

- Fever over 103 F (39.4 C)
- Chest pain or difficulty breathing occurs
- You become worse.

12 **Care Advice given per Coronavirus (COVID-19) - Diagnosed or Suspected (Adult) guideline.**

## Home Care

[1] COVID-19 diagnosed by positive lab test AND [2] mild symptoms (e.g., cough, fever, others) AND [3] no complications or SOB

1 **Home Care:**

- You should be able to treat this at home.

2 **Reassurance and Education - Positive COVID-19 Lab Test and Mild Symptoms:**

- You had a lab test for COVID-19 and it came back positive.
- The diagnosis has been confirmed.
- From what you have told me, your symptoms are mild. That is reassuring.
- *Here's some care advice to help you and to help prevent others from getting sick.*

3 **General Care Advice for COVID-19 Symptoms:**

- The treatment is the same whether you have COVID-19, influenza or some other respiratory virus.
- **Cough:** Use cough drops.
- **Feeling dehydrated:** Drink extra liquids. If the air in your home is dry, use a humidifier.
- **Fever:** For fever over 101 F (38.3 C), take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package. Do not take aspirin unless your doctor has prescribed it for you.
- **Muscle aches, headache, and other pains:** Often this comes and goes with the fever. Take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package.
- **Sore throat:** Try throat lozenges, hard candy or warm chicken broth.

4 **Cough Medicines:**

- **OTC Cough Syrups:** The most common cough suppressant in OTC cough medications is dextromethorphan. Often the letters "DM" appear in the name.
- **OTC Cough Drops:** Cough drops can help a lot, especially for mild coughs. They reduce coughing by soothing your irritated throat and removing that tickle sensation in the back of the throat. Cough drops also have the advantage of portability - you can carry them with you.
- **Home Remedy - Hard Candy:** Hard candy works just as well as medicine-flavored OTC cough drops. People who have diabetes should use sugar-free candy.
- **Home Remedy - Honey:** This old home remedy has been shown to help decrease coughing at night. The adult dosage is 2 teaspoons (10 ml) at bedtime. Honey should not be given to infants under one year of age.

**5 Caution - Dextromethorphan:**

- Do not try to completely suppress coughs that produce mucus and phlegm. Remember that coughing is helpful in bringing up mucus from the lungs and preventing pneumonia.
- Research Notes: Dextromethorphan in some research studies has been shown to reduce the frequency and severity of cough in adults (18 years or older) without significant adverse effects. However, other studies suggest that dextromethorphan is no better than placebo at reducing a cough.
- Drug Abuse Potential: It should be noted that dextromethorphan has become a drug of abuse. This problem is seen most often in adolescents. Overdose symptoms can range from giggling and euphoria to hallucinations and coma.
- **Contraindicated:** Do not take dextromethorphan if you are taking a monoamine oxidase (MAO) inhibitor now or in the past 2 weeks. Examples of MAO inhibitors include isocarboxazid (Marplan), phenelzine (Nardil), selegiline (Eldepryl), Emsam, Zelapar), and tranlycypromine (Parnate). Do not take dextromethorphan if you are taking venlafaxine (Effexor).

**6 Humidifier:**

- If the air is dry, use a humidifier in the bedroom.
- Dry air makes coughs worse.

**7 Coughing Spells:**

- Drink warm fluids. Inhale warm mist. (Reason: both relax the airway and loosen up the phlegm)
- Suck on cough drops or hard candy to coat the irritated throat.

**8 Avoid Tobacco Smoke:**

- Avoid tobacco smoke.
- Smoking or being exposed to smoke makes coughs much worse.

**9 Pain and Fever Medicines:**

- For pain or fever relief, take either acetaminophen or ibuprofen.
- They are over-the-counter (OTC) drugs that help treat both fever and pain. You can buy them at the drugstore.
- Treat fevers above 101° F (38.3° C). The goal of fever therapy is to bring the fever down to a comfortable level. Remember that fever medicine usually lowers fever 2 degrees F (1 - 1 1/2 degrees C).
- **Acetaminophen Regular Strength Tylenol:** Take 650 mg (two 325 mg pills) by mouth every 4 to 6 hours as needed. Each Regular Strength Tylenol pill has 325 mg of acetaminophen. The most you should take each day is 3,250 mg (10 pills a day).
- **Acetaminophen - Extra Strength Tylenol:** Take 1,000 mg (two 500 mg pills) every 8 hours as needed. Each Extra Strength Tylenol pill has 500 mg of acetaminophen. The most you should take each day is 3,000 mg (6 pills a day).
- **Ibuprofen (e.g., Motrin, Advil):** Take 400 mg (two 200 mg pills) by mouth every 6 hours. The most you should take each day is 1,200 mg (six 200 mg pills), unless your doctor has told you to take more.

**10 Pain and Fever Medicines - Extra Notes and Warnings:**

- Use the lowest amount of medicine that makes your pain or fever better.
- Acetaminophen is thought to be safer than ibuprofen or naproxen in people over 65 years old. Acetaminophen is in many OTC and prescription medicines. It might be in more than one medicine that you are taking. You need to be careful and not take an overdose. An acetaminophen overdose can hurt the liver.
- McNeil, the company that makes Tylenol, has different dosage instructions for Tylenol in Canada and the United States. In Canada, the maximum recommended dose per day is 4,000 mg or twelve Regular-Strength (325 mg) pills. In the United States, McNeil recommends a maximum dose of ten Regular-Strength (325 mg) pills.
- **Caution:** Do not take acetaminophen if you have liver disease.
- **Caution:** Do not take ibuprofen if you have stomach problems, kidney disease, are pregnant, or have been told by your doctor to avoid this type of anti-inflammatory drug. Do not take ibuprofen for more than 7 days without consulting your doctor.
- *Before taking any medicine, read all the instructions on the package.*

**11 No Aspirin:**

- Do not use aspirin for treatment of fever or pain.
- *Reason:* there is an association between influenza and Reyes' Syndrome.

**12 Note to Triager - Ibuprofen Concerns:**

- Discuss only if caller brings up concerns about ibuprofen.
- **Response:** The CDC, WHO, and other experts continue to support the use of ibuprofen (if needed) for patients with COVID-19. They found no scientific evidence to support the claim that ibuprofen made COVID-19 worse.

- 13 **How to Protect Others - When You are Sick with COVID-19:**
- **Stay Home a Minimum of 10 Days:** Home isolation is needed for at least 10 days after the symptoms started. Stay home from school or work if you are sick. Do **Not** go to religious services, child care centers, shopping, or other public places. Do **Not** use public transportation (e.g., bus, taxis, ride-sharing). Do **Not** allow any visitors to your home. Leave the house only if you need to seek urgent medical care.
  - **Cover the Cough:** Cough and sneeze into your shirt sleeve or inner elbow. Don't cough into your hand or the air. If available, cough into a tissue and throw it into a trash can.
  - **Wash Hands Often:** Wash hands often with soap and water. After coughing or sneezing are important times. If soap and water are not available, use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry. Avoid touching your eyes, nose, and mouth with unwashed hands.
  - **Wear a Mask:** Wear a facemask when around others. Always wear a facemask (if available) if you have to leave your home (such as going to a medical facility).
  - **Call First if Medical Care Needed:** Call ahead to get approval and careful directions.
- 14 **How and When to Stop Home Isolation:**
- *Talk to your healthcare provider.*
  - The decision to stop home isolation should be made by your healthcare provider in consultation with the local health departments. Local decisions depend on local circumstances.
- 15 **Stopping Home Isolation - Must Meet all 3 Requirements (CDC):**
- Fever gone for at least 24 hours off fever-reducing medicines **AND**
  - Cough and other symptoms must be improved **AND**
  - Symptoms started more than 10 days ago.
  - *If unsure if it is safe for you to leave isolation, check the CDC website or call your healthcare provider.*
- 16 **Other COVID-19 Facts:**
- **Incubation Period:** Average 5 days (range 2 to 14 days) after coming in contact with a person who has COVID-19 virus.
  - **No Symptoms, but Infected (Asymptomatic):** Approximately 30% of infected patients may have no symptoms.
  - **Mild Infections:** About 80% of those with symptoms have a mild illness, much like a normal flu or a bad cold. The symptoms usually last 2 weeks.
  - **Severe Infections:** About 20% of those with symptoms develop trouble breathing from viral pneumonia. Many of these need to be admitted to the hospital. People with complications generally recover in 3 to 6 weeks.
  - **Death Rate:** The adult death rate is approximately 1% to 3%. The death rate is lower in children and younger adults. It is higher in older adults.
  - **Treatment:** Treatment is supportive. Oxygen and IV fluids are used for hospitalized patients. On May 1, 2020 the U.S. Food and Drug Administration (FDA) granted temporary emergency use authorization (EUA) for the investigational antiviral remdesivir to treat COVID-19. Doctors can use this drug to treat patients who are hospitalized with COVID-19 who are severely ill. More research is needed to determine how well this drug works and for which patients.
  - **Prevention - Vaccine:** There currently is no approved vaccine to prevent COVID-19. There are vaccines in development. A COVID-19 vaccine may be available for adults by early 2021.
  - **Prevention - Medicine:** Currently, there is no medicine to prevent COVID-19. Warning: the malaria drug (chloroquine) was studied and found not to be helpful for this disease. It also had cardiac side effects. *Social distancing and wearing masks have been proven to help prevent COVID-19!*
- 17 **Call Back If:**
- Fever over 103 F (39.4 C)
  - Fever lasts over 3 days
  - Fever returns after being gone for 24 hours
  - Chest pain or difficulty breathing occurs
  - You become worse.
- 18 **Care Advice** given per **Coronavirus (COVID-19) - Diagnosed or Suspected** (Adult) guideline.

[1] COVID-19 diagnosed by HCP (doctor, NP or PA) AND [2] mild symptoms (e.g., cough, fever, others) AND [3] no complications or SOB

- 1 **Home Care:**
- You should be able to treat this at home.

- 2 Reassurance and Education - Diagnosed with COVID-19 by Health Care Provider (HCP) and Mild Symptoms:**
- Your HCP has diagnosed you as having COVID-19 based on your symptoms and other information.
  - You may or may not have not been tested for COVID-19. People with mild symptoms sometimes do not need testing. It does not change treatment.
  - The symptoms of COVID-19 can be mild, especially if you are healthy and under 65 years old.
  - *Here's some care advice to help you and to help prevent others from getting sick.*
- 3 General Care Advice for COVID-19 Symptoms:**
- The treatment is the same whether you have COVID-19, influenza or some other respiratory virus.
  - **Cough:** Use cough drops.
  - **Feeling dehydrated:** Drink extra liquids. If the air in your home is dry, use a humidifier.
  - **Fever:** For fever over 101 F (38.3 C), take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package. Do not take aspirin unless your doctor has prescribed it for you.
  - **Muscle aches, headache, and other pains:** Often this comes and goes with the fever. Take acetaminophen every 4-6 hours (Adults 650 mg) OR ibuprofen every 6-8 hours (Adults 400 mg). Before taking any medicine, read all the instructions on the package.
  - **Sore throat:** Try throat lozenges, hard candy or warm chicken broth.
- 4 Cough Medicines:**
- **OTC Cough Syrups:** The most common cough suppressant in OTC cough medications is dextromethorphan. Often the letters "DM" appear in the name.
  - **OTC Cough Drops:** Cough drops can help a lot, especially for mild coughs. They reduce coughing by soothing your irritated throat and removing that tickle sensation in the back of the throat. Cough drops also have the advantage of portability - you can carry them with you.
  - **Home Remedy - Hard Candy:** Hard candy works just as well as medicine-flavored OTC cough drops. People who have diabetes should use sugar-free candy.
  - **Home Remedy - Honey:** This old home remedy has been shown to help decrease coughing at night. The adult dosage is 2 teaspoons (10 ml) at bedtime. Honey should not be given to infants under one year of age.
- 5 Caution - Dextromethorphan:**
- Do not try to completely suppress coughs that produce mucus and phlegm. Remember that coughing is helpful in bringing up mucus from the lungs and preventing pneumonia.
  - **Research Notes:** Dextromethorphan in some research studies has been shown to reduce the frequency and severity of cough in adults (18 years or older) without significant adverse effects. However, other studies suggest that dextromethorphan is no better than placebo at reducing a cough.
  - **Drug Abuse Potential:** It should be noted that dextromethorphan has become a drug of abuse. This problem is seen most often in adolescents. Overdose symptoms can range from giggling and euphoria to hallucinations and coma.
  - **Contraindicated:** Do not take dextromethorphan if you are taking a monoamine oxidase (MAO) inhibitor now or in the past 2 weeks. Examples of MAO inhibitors include isocarboxazid (Marplan), phenelzine (Nardil), selegiline (Eldepryl, Emsam, Zelapar), and tranylcypromine (Parnate). Do not take dextromethorphan if you are taking venlafaxine (Effexor).
- 6 Humidifier:**
- If the air is dry, use a humidifier in the bedroom.
  - Dry air makes coughs worse.
- 7 Coughing Spells:**
- Drink warm fluids. Inhale warm mist. (Reason: both relax the airway and loosen up the phlegm)
  - Suck on cough drops or hard candy to coat the irritated throat.
- 8 Avoid Tobacco Smoke:**
- Avoid tobacco smoke.
  - Smoking or being exposed to smoke makes coughs much worse.
- 9 Pain and Fever Medicines:**
- For pain or fever relief, take either acetaminophen or ibuprofen.
  - They are over-the-counter (OTC) drugs that help treat both fever and pain. You can buy them at the drugstore.
  - Treat fevers above 101° F (38.3° C). The goal of fever therapy is to bring the fever down to a comfortable level. Remember that fever medicine usually lowers fever 2 degrees F (1 - 1 1/2 degrees C).
  - **Acetaminophen Regular Strength Tylenol:** Take 650 mg (two 325 mg pills) by mouth every 4 to 6 hours as needed. Each Regular Strength Tylenol pill has 325 mg of acetaminophen. The most you should take each day is 3,250 mg (10 pills a day).
  - **Acetaminophen - Extra Strength Tylenol:** Take 1,000 mg (two 500 mg pills) every 8 hours as needed. Each Extra Strength Tylenol pill has 500 mg of acetaminophen. The most you should take each day is 3,000 mg (6 pills a day).
  - **Ibuprofen (e.g., Motrin, Advil):** Take 400 mg (two 200 mg pills) by mouth every 6 hours. The most you should take each day is 1,200 mg (six 200 mg pills), unless your doctor has told you to take more.

- 10 **Pain and Fever Medicines - Extra Notes and Warnings:**
- Use the lowest amount of medicine that makes your pain or fever better.
  - Acetaminophen is thought to be safer than ibuprofen or naproxen in people over 65 years old. Acetaminophen is in many OTC and prescription medicines. It might be in more than one medicine that you are taking. You need to be careful and not take an overdose. An acetaminophen overdose can hurt the liver.
  - McNeil, the company that makes Tylenol, has different dosage instructions for Tylenol in Canada and the United States. In Canada, the maximum recommended dose per day is 4,000 mg or twelve Regular-Strength (325 mg) pills. In the United States, McNeil recommends a maximum dose of ten Regular-Strength (325 mg) pills.
  - **Caution:** Do not take acetaminophen if you have liver disease.
  - **Caution:** Do not take ibuprofen if you have stomach problems, kidney disease, are pregnant, or have been told by your doctor to avoid this type of anti-inflammatory drug. Do not take ibuprofen for more than 7 days without consulting your doctor.
  - *Before taking any medicine, read all the instructions on the package.*
- 11 **No Aspirin:**
- Do not use aspirin for treatment of fever or pain.
  - *Reason:* there is an association between influenza and Reyes' Syndrome.
- 12 **Note to Triager - Ibuprofen Concerns:**
- Discuss only if caller brings up concerns about ibuprofen.
  - **Response:** The CDC, WHO, and other experts continue to support the use of ibuprofen (if needed) for patients with COVID-19. They found no scientific evidence to support the claim that ibuprofen made COVID-19 worse.
- 13 **How to Protect Others - When You are Sick with COVID-19:**
- **Stay Home a Minimum of 10 Days:** Home isolation is needed for at least 10 days after the symptoms started. Stay home from school or work if you are sick. Do **Not** go to religious services, child care centers, shopping, or other public places. Do **Not** use public transportation (e.g., bus, taxis, ride-sharing). Do **Not** allow any visitors to your home. Leave the house only if you need to seek urgent medical care.
  - **Cover the Cough:** Cough and sneeze into your shirt sleeve or inner elbow. Don't cough into your hand or the air. If available, cough into a tissue and throw it into a trash can.
  - **Wash Hands Often:** Wash hands often with soap and water. After coughing or sneezing are important times. If soap and water are not available, use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry. Avoid touching your eyes, nose, and mouth with unwashed hands.
  - **Wear a Mask:** Wear a facemask when around others. Always wear a facemask (if available) if you have to leave your home (such as going to a medical facility).
  - **Call First if Medical Care Needed:** Call ahead to get approval and careful directions.
- 14 **COVID-19 - Face Masks for Prevention:**
- Face masks are essential for reducing the spread of COVID-19. They will also reduce the spread of influenza. Wearing a mask means you care about other people.
  - **Sick Patients:** People who are sick with COVID-19 must always wear a face mask if they need to leave the home. Example: for medical visits. *Exception:* patients with trouble breathing (CDC). Consider a loose face covering such as a bandana.
  - **Well People:** When community spread is high, the CDC also recommends face masks or coverings for everyone going outside the home. They are critical if entering a public building, such as a grocery store. *Reason:* Many people with COVID-19 have no symptoms but can spread the virus. *Exceptions:* Face mask or covering is optional if outdoors and you can avoid being within 6 feet (2 meters) of other people. Some examples are an outdoor walk or run.
  - **Age Limits:** Face coverings also are not recommended for children under 2 years (CDC).
  - More information on how to make and use a cloth face mask at: <https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/how-to-make-cloth-face-covering.html>.
- 15 **How and When to Stop Home Isolation:**
- *Talk to your healthcare provider.*
  - The decision to stop home isolation should be made by your healthcare provider in consultation with the local health departments. Local decisions depend on local circumstances.
- 16 **Stopping Home Isolation - Must Meet all 3 Requirements (CDC):**
- Fever gone for at least 24 hours off fever-reducing medicines **AND**
  - Cough and other symptoms must be improved **AND**
  - Symptoms started more than 10 days ago.
  - *If unsure if it is safe for you to leave isolation, check the CDC website or call your healthcare provider.*



17 **Other COVID-19 Facts:**

- **Incubation Period:** Average 5 days (range 2 to 14 days) after coming in contact with a person who has COVID-19 virus.
- **No Symptoms, but Infected (Asymptomatic):** Approximately 30% of infected patients may have no symptoms.
- **Mild Infections:** About 80% of those with symptoms have a mild illness, much like a normal flu or a bad cold. The symptoms usually last 2 weeks.
- **Severe Infections:** About 20% of those with symptoms develop trouble breathing from viral pneumonia. Many of these need to be admitted to the hospital. People with complications generally recover in 3 to 6 weeks.
- **Death Rate:** The adult death rate is approximately 1% to 3%. The death rate is lower in children and younger adults. It is higher in older adults.
- **Treatment:** Treatment is supportive. Oxygen and IV fluids are used for hospitalized patients. On May 1, 2020 the U.S. Food and Drug Administration (FDA) granted temporary emergency use authorization (EUA) for the investigational antiviral remdesivir to treat COVID-19. Doctors can use this drug to treat patients who are hospitalized with COVID-19 who are severely ill. More research is needed to determine how well this drug works and for which patients.
- **Prevention - Vaccine:** There currently is no approved vaccine to prevent COVID-19. There are vaccines in development. A COVID-19 vaccine may be available for adults by early 2021.
- **Prevention - Medicine:** Currently, there is no medicine to prevent COVID-19. Warning: the malaria drug (chloroquine) was studied and found not to be helpful for this disease. It also had cardiac side effects. *Social distancing and wearing masks have been proven to help prevent COVID-19!*

18 **Call Back If:**

- Fever over 103 F (39.4 C)
- Fever lasts over 3 days
- Fever returns after being gone for 24 hours
- Chest pain or difficulty breathing occurs
- You become worse.

19 **Care Advice** given per **Coronavirus (COVID-19) - Diagnosed or Suspected (Adult)** guideline.

COVID-19 Home Isolation, questions about

*Note: How to protect others when you are sick.*

1 **Home Care:**

- You should be able to treat this at home.

2 **How to Protect Others - When You are Sick with COVID-19:**

- **Stay Home a Minimum of 10 Days:** Home isolation is needed for at least 10 days after the symptoms started. Stay home from school or work if you are sick. Do **Not** go to religious services, child care centers, shopping, or other public places. Do **Not** use public transportation (e.g., bus, taxis, ride-sharing). Do **Not** allow any visitors to your home. Leave the house only if you need to seek urgent medical care.
- **Cover the Cough:** Cough and sneeze into your shirt sleeve or inner elbow. Don't cough into your hand or the air. If available, cough into a tissue and throw it into a trash can.
- **Wash Hands Often:** Wash hands often with soap and water. After coughing or sneezing are important times. If soap and water are not available, use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry. Avoid touching your eyes, nose, and mouth with unwashed hands.
- **Wear a Mask:** Wear a facemask when around others. Always wear a facemask (if available) if you have to leave your home (such as going to a medical facility).
- **Call First if Medical Care Needed:** Call ahead to get approval and careful directions.

3 **Clean Your Hands Often:**

- **Wash Hands:** Wash your hands often with soap and water for at least 20 seconds. This is especially important after blowing your nose, coughing, or sneezing; going to the bathroom; and before eating or preparing food.
- **Use Hand Sanitizer:** If soap and water are not available, use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry.
- Avoid touching your eyes, nose, and mouth with unwashed hands.

4 **Clean "High Touch" Surfaces Every Day:**

- Clean high-touch surfaces in your isolation area ("sick room" and bathroom) every day.
- High-touch surfaces include phones, remote controls, counters, tabletops, doorknobs, bathroom fixtures, toilets, keyboards, tablets, and bedside tables.

5 **Stay Away From Others in Your Home:**

- If possible, stay in a specific "sick room" and away from other people in your home.
- Use a separate bathroom, if available.

- 6 **Call Ahead Before Visiting Your Healthcare Provider (Doctor, NP, PA):**
  - Call ahead: If you have a medical appointment, call your doctor's office or emergency department, and tell them you have or may have COVID-19.
  - This will help the office protect themselves and other patients.
  - Wear a facemask.
- 7 **How and When to Stop Home Isolation:**
  - *Talk to your healthcare provider.*
  - The decision to stop home isolation should be made by your healthcare provider in consultation with the local health departments. Local decisions depend on local circumstances.
- 8 **Stopping Home Isolation - Must Meet all 3 Requirements (CDC):**
  - Fever gone for at least 24 hours off fever-reducing medicines **AND**
  - Cough and other symptoms must be improved **AND**
  - Symptoms started more than 10 days ago.
  - *If unsure if it is safe for you to leave isolation, check the CDC website or call your healthcare provider.*
- 9 **Call Back If:**
  - You have more questions.
- 10 **Care Advice given per Coronavirus (COVID-19) - Diagnosed or Suspected (Adult) guideline.**

## COVID-19 Testing, questions about

- 1 **Home Care:**
  - You should be able to treat this at home.
- 2 **Note to Triager - COVID-19 Testing:**
  - **For questions about testing, it is best to direct the patient to their HCP** (doctor, NP, PA) during office hours. The HCP is the best resource for up-to-date information on testing. This is a somewhat complicated decision process. National and state testing recommendations continue to change. *Testing requires a HCP's order (as with all medical tests).*
  - COVID-19 testing is becoming more available from local and state public health departments. Commercial labs now can perform COVID-19 testing. However, it may still be difficult to find a place to get tested.
  - **What about a standing order?** As testing becomes more widely available, call centers should talk with their medical leadership about triagers being able to use a standing order for testing.
- 3 **COVID-19 - Who Needs Testing?**
  - People who have **symptoms of COVID-19**.
  - People who have had **close contact**, within 6 feet (2 meters) for a total of 15 minutes or more in a 24-hour period, with someone with confirmed to have COVID-19. People who **live with someone** confirmed to have COVID-19.
  - People living or working in a congregate setting (such as a correctional facility, long-term care facility, or shelter) where an **outbreak has occurred**.
- 4 **COVID-19 - Where to Go for Testing?**
  - Go to the testing site recommended by your healthcare provider (e.g., doctor, NP, or PA) or public health department.
  - Swabs of the nose or throat will only be collected on patients who have a healthcare provider's order.
  - Testing sites vary based on the city, hospital, and healthcare system.
  - In general, they are not performed in private doctor's offices or clinics.
  - *People cannot just walk in and request a COVID-19 test.*
- 5 **COVID-19 - What Types of Tests Are Available?**
  - There are two types of tests available for COVID-19: viral tests and antibody tests.
  - **Test for Current Infection - Viral Test:** A viral test tells you if you have the COVID-19 infection right now. A viral test is done with either a nasal swab or a saliva sample. Symptomatic patients should get a test within 3 days, if available. Asymptomatic people with a close contact COVID-19 exposure should get the viral test on day 6 to 8 after exposure.
  - **Test for Past Infection - Antibody Test:** An antibody test tells you if you have had COVID-19 before. This test is done with a blood sample. An antibody test may not be able to show if you have a current infection, because it can take 1 to 3 weeks for your body to make antibodies to the infection. We do not know yet if having antibodies to the virus can protect someone from getting infected with the virus again, or how long that protection might last. If an antibody test is needed, it is usually performed 2 to 3 weeks after the start of the infection.
  - The results usually come back in 1 to 3 days, but may take longer depending on testing kit or testing site availability.

## 6 Understanding Viral Test Results:

- Viral tests look for part of the virus (RNA or proteins) in the test sample. Since this test looks for actual parts of the virus, it can tell you if you are *currently infected* (active infection) and at risk of spreading the disease. A viral test is done using a nasal (nose) swab or saliva (spit).
- A **positive viral test** means that you most likely have an active COVID-19 infection and *can spread the infection to others*.
- A **negative viral test** means that you likely did NOT have an active COVID-19 infection at the time the test was done.
- *Note:* All tests can sometimes have a false (wrong) result. Some reasons for this include how the sample was collected, how long into the illness it was taken, and the type of test done. That is why it is important to discuss your results with your health care provider.

## 7 Understanding Antibody Test Results:

- Antibody tests (also called serology tests) are done on blood samples.
- COVID-19 antibody tests look for antibodies to the virus that causes COVID-19. Antibodies begin to form during an infection but can last as long as a lifetime. Therefore, an antibody test cannot tell the difference between an active infection and one you had sometime in the past.
- It is not yet known if antibodies protect you from getting COVID-19 again.
- A **positive antibody test** means that you most likely *have been*, or *are now*, infected with COVID-19.
- A **negative antibody test** means that you likely *never had* COVID 19 OR you *have not yet* formed antibodies to COVID-19.

## 8 Understanding Viral and Antibody Testing Together:

- In some cases, your healthcare provider (HCP) may want to perform both antibody testing and viral testing together. The best source of information on understanding the test results will come from your HCP. Here is some information on how the two results can be used together.
- **Both Tests Are Positive:** You most likely have an *active infection* and can spread COVID-19 to others.
- **Both Tests Are Negative:** You likely do NOT have COVID-19 and likely never had a COVID-19 infection.
- **Viral Test Is Positive and Antibody Test Is Negative:** You most likely have an *active infection* and can spread COVID-19 to others. You have not yet developed antibodies to your active COVID-19 infection.
- **Viral Test Is Negative and Antibody Test Is Positive:** You likely do NOT have an active COVID-19 infection but did have COVID-19 sometime in the past.

## 9 Call Back If:

- You have more questions.

## 10 Care Advice given per Coronavirus (COVID-19) - Diagnosed or Suspected (Adult) guideline.

### COVID-19 Prevention and Healthy Living, questions about

*Note: How to protect you and your family; how to reduce anxiety and stress.*

#### 1 Home Care:

- You should be able to treat this at home.

#### 2 COVID-19 - Face Masks for Prevention:

- Face masks are essential for reducing the spread of COVID-19. They will also reduce the spread of influenza. Wearing a mask means you care about other people.
- **Sick Patients:** People who are sick with COVID-19 must always wear a face mask if they need to leave the home. Example: for medical visits. *Exception:* patients with trouble breathing (CDC). Consider a loose face covering such as a bandana.
- **Well People:** When community spread is high, the CDC also recommends face masks or coverings for everyone going outside the home. They are critical if entering a public building, such as a grocery store. *Reason:* Many people with COVID-19 have no symptoms but can spread the virus. *Exceptions:* Face mask or covering is optional if outdoors and you can avoid being within 6 feet (2 meters) of other people. Some examples are an outdoor walk or run.
- **Age Limits:** Face coverings also are not recommended for children under 2 years (CDC).
- More information on how to make and use a cloth face mask at: <https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/how-to-make-cloth-face-covering.html>.

- 3 **COVID-19 - How to Protect Your Family and Yourself From Getting Sick:**
  - Avoid close contact with people known to have this new coronavirus infection.
  - Avoid close contact with people outside your family unit.
  - Avoid closed spaces (indoors) when possible and all crowds (even outdoors).
  - When you must leave your home, wear a mask and observe social (safe) distancing.
  - Try to stay at least 6 feet (2 meters) away from anyone who is coughing.
  - Wash hands often with soap and water.
  - Alcohol-based hand cleaners are also effective.
  - Avoid touching the eyes, nose or mouth. Germs on the hands can spread this way.
  - Do not share eating utensils (e.g., spoon, fork).
- 4 **Keep Your Mind Positive:**
  - **Live in the Present:** Live in the present, not the future. The future is where your needless worries live.
  - **Think Positive:** Use a mantra to reduce your fears, such as "I am strong". Stay positive.
  - **Get Outdoors:** Take daily walks. Go to a park if you have one. Being in nature is good for your immune system.
  - **Stay in Touch With Your Friends and Family:** Use regular phone calls and video chats to stay in touch with those you love. Schedule virtual video dinners with friends and family!
- 5 **Keep Your Body Strong:**
  - Get your body ready to fight the COVID-19 virus.
  - Get enough sleep.
  - Stay physically active. Walk or exercise every day. Take the stairs.
  - Stay well hydrated.
  - Eat healthy meals. Avoid overeating to deal with your fears.
  - Avoid the over-use of anti-fever medicines. Fever helps fight infections and ramps up your immune system.
- 6 **Ask for Help:**
  - If you feel so sad or worried that you can not function, reach out to your health care provider, local mental health center, or SAMHSA's National Helpline.
  - **SAMHSA:** 1-800-662-HELP (4357). Website: [www.samhsa.gov/find-help/national-helpline](http://www.samhsa.gov/find-help/national-helpline).
- 7 **Get a Flu Shot (Influenza Vaccine):**
  - Protect yourself and your family from influenza by getting your annual flu shot (influenza vaccination).
  - All adults should get a flu shot every year. This year is more important than ever. *Reason:* Getting COVID-19 while you also have or are recovering from the flu may increase the chances of getting severe symptoms.
- 8 **How to Protect Others - When You are Sick with COVID-19:**
  - **Stay Home a Minimum of 10 Days:** Home isolation is needed for at least 10 days after the symptoms started. Stay home from school or work if you are sick. Do **Not** go to religious services, child care centers, shopping, or other public places. Do **Not** use public transportation (e.g., bus, taxis, ride-sharing). Do **Not** allow any visitors to your home. Leave the house only if you need to seek urgent medical care.
  - **Cover the Cough:** Cough and sneeze into your shirt sleeve or inner elbow. Don't cough into your hand or the air. If available, cough into a tissue and throw it into a trash can.
  - **Wash Hands Often:** Wash hands often with soap and water. After coughing or sneezing are important times. If soap and water are not available, use an alcohol-based hand sanitizer with at least 60% alcohol, covering all surfaces of your hands and rubbing them together until they feel dry. Avoid touching your eyes, nose, and mouth with unwashed hands.
  - **Wear a Mask:** Wear a facemask when around others. Always wear a facemask (if available) if you have to leave your home (such as going to a medical facility).
  - **Call First if Medical Care Needed:** Call ahead to get approval and careful directions.
- 9 **Call Back If:**
  - You have more questions.
- 10 **Care Advice** given per **Coronavirus (COVID-19) - Diagnosed or Suspected** (Adult) guideline.

## COVID-19 Disease, questions about

*Note: Broad information including symptoms, how it is spread, travel, et.al.*

- 1 **Home Care:**
  - You should be able to treat this at home.

- 2 **COVID-19 (Coronavirus Disease 2019) - Introduction:**
  - An outbreak of this infection began in China in December 2019.
  - The first patient in the United States occurred on January 21, 2020.
  - Four patients were confirmed in Canada on January 31, 2020.
  - The *World Health Organization* (WHO) declared COVID-19 a global public health emergency on January 30, 2020 and then a pandemic on March 11, 2020.
  - The *Centers for Disease Control and Prevention* (CDC) is considered the source of truth for this guideline. This continues to be a rapidly changing situation and guidance from the CDC is being updated daily. See <https://www.cdc.gov/coronavirus/2019-nCoV/index.html>.
- 3 **COVID-19 - Symptoms:**
  - COVID-19 most often causes a respiratory illness.
  - *The most common symptoms are:* cough, fever, and shortness of breath.
  - *Other less common symptoms are:* chills, fatigue, headache, loss of smell or taste, muscle pain, and sore throat.
  - Some people may have minimal symptoms or even have no symptoms (asymptomatic).
- 4 **COVID-19 - Exposure Risk Factors:**
  - *Here are the main risk factors for getting sick with COVID-19.*
  - **Close contact with a person** who tested positive for COVID-19 AND contact occurred while they were ill. Close contact means being within 6 feet (2 meters) for a total of 15 minutes or more in a 24-hour period. This includes living with someone infected with COVID-19.
  - **Living in or travel from a city** or area where there is community spread of COVID-19. This carries a lower risk compared to close contact if one follows physical distancing recommendations. Community spread has occurred in most of the US, especially in cities.
  - **International Travel:** The CDC (<https://www.cdc.gov/coronavirus/2019-ncov/travelers>) has the most up-to-date list of where COVID-19 outbreaks are occurring.
- 5 **COVID-19 - How It Is Spread:**
  - *COVID-19 is spread from person to person.*
  - **Respiratory Droplets:** The virus spreads from respiratory droplets that are produced when a person coughs, sneezes, shouts, or sings. The infected droplets can then be inhaled by a nearby person or land on the surface of their face or eyes. Droplets fall quickly to the floor or ground. This is **how most COVID is spread**.
  - **Surfaces:** Most infected people also have respiratory secretions on their hands. These secretions get transferred to healthy people on doorknobs, faucet handles, etc. The virus then gets transferred to healthy people when they touch their face or rub their eyes. This is a **less common cause** of spread.
  - **Aerosols:** Aerosols are tiny, invisible particles that can float in the air for 1 to 2 hours and travel more than 6 feet (2 meters). They only occur in a closed room with poor ventilation. Aerosols are a **rare cause** of COVID-19 spread.
- 6 **COVID-19 - How to Protect Your Family and Yourself From Getting Sick:**
  - *Avoid close contact with people known to have this new coronavirus infection.*
  - *Avoid close contact with people outside your family unit.*
  - Avoid closed spaces (indoors) when possible and all crowds (even outdoors).
  - When you must leave your home, wear a mask and observe social (safe) distancing.
  - Try to stay at least 6 feet (2 meters) away from anyone who is coughing.
  - Wash hands often with soap and water.
  - Alcohol-based hand cleaners are also effective.
  - Avoid touching the eyes, nose or mouth. Germs on the hands can spread this way.
  - Do not share eating utensils (e.g., spoon, fork).
- 7 **COVID-19 - Travel Guidelines:**
  - The Centers for Disease Control and Prevention (CDC) maintains a website with latest recommendations regarding travel and your health.
  - Currently the CDC recommends against travel to many geographic areas with widespread and ongoing spread of COVID-19. See current list at <https://wwwnc.cdc.gov/travel/>.
  - *CDC Travel Health Website:* <https://wwwnc.cdc.gov/travel/>.
  - *CDC Travel FAQs:* <https://www.cdc.gov/coronavirus/2019-ncov/travelers/faqs.html>

## 8 COVID-19 - What Types of Tests Are Available?

- There are two types of tests available for COVID-19: viral tests and antibody tests.
- **Test for Current Infection - Viral Test:** A viral test tells you if you have the COVID-19 infection right now. A viral test is done with either a nasal swab or a saliva sample. Symptomatic patients should get a test within 3 days, if available. Asymptomatic people with a close contact COVID-19 exposure should get the viral test on day 6 to 8 after exposure.
- **Test for Past Infection - Antibody Test:** An antibody test tells you if you have had COVID-19 before. This test is done with a blood sample. An antibody test may not be able to show if you have a current infection, because it can take 1 to 3 weeks for your body to make antibodies to the infection. We do not know yet if having antibodies to the virus can protect someone from getting infected with the virus again, or how long that protection might last. If an antibody test is needed, it is usually performed 2 to 3 weeks after the start of the infection.
- The results usually come back in 1 to 3 days, but may take longer depending on testing kit or testing site availability.

## 9 COVID-19 - Face Masks for Prevention:

- Face masks are essential for reducing the spread of COVID-19. They will also reduce the spread of influenza. Wearing a mask means you care about other people.
- **Sick Patients:** People who are sick with COVID-19 must always wear a face mask if they need to leave the home. Example: for medical visits. *Exception:* patients with trouble breathing (CDC). Consider a loose face covering such as a bandana.
- **Well People:** When community spread is high, the CDC also recommends face masks or coverings for everyone going outside the home. They are critical if entering a public building, such as a grocery store. *Reason:* Many people with COVID-19 have no symptoms but can spread the virus. *Exceptions:* Face mask or covering is optional if outdoors and you can avoid being within 6 feet (2 meters) of other people. Some examples are an outdoor walk or run.
- **Age Limits:** Face coverings also are not recommended for children under 2 years (CDC).
- More information on how to make and use a cloth face mask at: <https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/how-to-make-cloth-face-covering.html>.

## 10 Reassurance and Education - Going to the ED or Urgent Care Center During the COVID-19 Pandemic:

- If you or your child needs to be seen for an urgent medical problem, do not hesitate to go.
- ERs and urgent care centers are safe places. They are well equipped to protect you against the virus.
- For non-urgent conditions, talk to your healthcare provider's office first.

## 11 Other COVID-19 Facts:

- **Incubation Period:** Average 5 days (range 2 to 14 days) after coming in contact with a person who has COVID-19 virus.
- **No Symptoms, but Infected (Asymptomatic):** Approximately 30% of infected patients may have no symptoms.
- **Mild Infections:** About 80% of those with symptoms have a mild illness, much like a normal flu or a bad cold. The symptoms usually last 2 weeks.
- **Severe Infections:** About 20% of those with symptoms develop trouble breathing from viral pneumonia. Many of these need to be admitted to the hospital. People with complications generally recover in 3 to 6 weeks.
- **Death Rate:** The adult death rate is approximately 1% to 3%. The death rate is lower in children and younger adults. It is higher in older adults.
- **Treatment:** Treatment is supportive. Oxygen and IV fluids are used for hospitalized patients. On May 1, 2020 the U.S. Food and Drug Administration (FDA) granted temporary emergency use authorization (EUA) for the investigational antiviral remdesivir to treat COVID-19. Doctors can use this drug to treat patients who are hospitalized with COVID-19 who are severely ill. More research is needed to determine how well this drug works and for which patients.
- **Prevention - Vaccine:** There currently is no approved vaccine to prevent COVID-19. There are vaccines in development. A COVID-19 vaccine may be available for adults by early 2021.
- **Prevention - Medicine:** Currently, there is no medicine to prevent COVID-19. Warning: the malaria drug (chloroquine) was studied and found not to be helpful for this disease. It also had cardiac side effects. *Social distancing and wearing masks have been proven to help prevent COVID-19!*

## 12 Call Back If:

- You have more questions.

## 13 Care Advice given per Coronavirus (COVID-19) - Diagnosed or Suspected (Adult) guideline.

## FIRST AID

N/A



## BACKGROUND INFORMATION

### Key Points



- An outbreak of this infection began in Wuhan, Hubei Province, China in December 2019.
- The first patient in the United States occurred on January 21, 2020. During March 2020 cases were identified in all states.
- Four patients were confirmed in Canada on January 31, 2020.
- The *World Health Organization* (WHO) declared COVID-19 a global public health emergency on January 30, 2020 and then a pandemic on March 11, 2020.
- The *Centers for Disease Control and Prevention* (CDC) is considered the source of truth for this guideline. This continues to be a rapidly changing situation and guidance from the CDC is being updated daily. See <https://www.cdc.gov/coronavirus/2019-nCoV/index.html>.

## Symptoms

The COVID-19 coronavirus most often causes a lower respiratory tract illness. More common symptoms are:

- Anorexia (40-84%)
- Chills (16-28%)
- **Cough** (59-82%)
- Fatigue (44-70%)
- **Fever** (83-99%)
- Loss of smell or taste (15%)
- Muscle pain (11-35%)
- **Shortness of breath or difficulty breathing** (31-40%)

Other symptoms may include:

- Diarrhea (2-5%)
- Headache (5-14%)
- Nausea or Vomiting (1-10%)
- Runny or Stuffy Nose (4%)
- Sore Throat (5%)
- Sputum production (14-34%)

Having two or more of any of these symptoms increases the likelihood that a person may have COVID-19.

Over 30% of infected adult patients have no symptoms (asymptomatic). Children and teens are even more likely to have no symptoms.

## Cause

It is caused by a novel (new) coronavirus (COVID-19).

## Exposure Risk Factors

Risk factors for getting sick with COVID-19 are:

- **Close contact with a person** who tested positive for COVID-19 AND contact occurred while they were ill. Close contact is defined as being within 6 feet (2 meters) of an infected person for a total of 15 minutes or more in a 24-hour period. This includes **living with someone** infected with COVID-19.
- **Living in or travel from a city** or area where there is community spread of COVID-19. This carries a lower risk compared to close contact if one follows physical distancing recommendations. Community spread is now occurring in most of the US, especially in cities.

• **International Travel:** The CDC (<https://www.cdc.gov/coronavirus/2019-ncov/travelers>) has the most up-to-date list of where COVID-19 outbreaks are occurring.

### How it is Spread (Transmission)

COVID-19 is spread from person to person.

• **Respiratory Droplets:** The virus spreads from respiratory droplets that are produced when a person coughs, sneezes, shouts, or sings. The infected droplets can then be inhaled by a nearby person or land on the surface of their face or eyes. Droplets fall quickly to the floor or ground. This is **how most COVID is spread**.

• **Surfaces:** Most infected people also have respiratory secretions on their hands. These secretions get transferred to healthy people on doorknobs, faucet handles, etc. The virus then gets transferred to healthy people when they touch their face or rub their eyes. This is a **less common cause** of spread.

• **Aerosols:** Aerosols are tiny, invisible particles that can float in the air for 1 to 2 hours and travel more than 6 feet (2 meters). They only occur in a closed room with poor ventilation. Aerosols are a **rare cause** of COVID-19 spread.

### Incubation Period

The incubation period averages 5 days (range 2 to 14 days) after coming in contact with the secretions of a person who has COVID-19.

### Testing

There are two types of tests for COVID-19: viral tests and antibody tests.

• **Test for Current Infection - Viral Test:** A viral test tells you if you have the COVID-19 infection right now. A viral test is done with either a nasal swab or a saliva sample.

• **Test for Past Infection - Antibody Test:** An antibody test tells you if you have had COVID-19 before. This test is done with a blood sample. An antibody test may not be able to show if you have a current infection, because it can take 1 to 3 weeks for your body to make antibodies to the infection. We do not know yet if having antibodies to the virus can protect someone from getting infected with the virus again, or how long that protection might last.

The results usually come back in 1 to 4 days, but may take longer depending on testing kit or testing site availability.

When should testing be performed?

• **People Who Have Symptoms of COVID-19:** A person who is symptomatic should get a COVID-19 viral test **within 3 days**.

• **People With Close Contact Exposure:** A person who had a close contact COVID-19 exposure and is asymptomatic should get a COVID-19 viral test **about 6 to 8 days** after exposure.

For questions about testing, it is often best to direct the patient to their HCP (doctor, NP, PA) during office hours. The HCP is the best resource for up-to-date information on testing. National and state testing recommendations continue to change. *Testing requires a HCP's order (as with all medical tests).*

### Diagnosis and Reporting

Clinicians should report positive test results to infection control at their health system and to their local or state health department.

## Complications

Complications include pneumonia, hypoxia, ARDS, respiratory failure, and death.

### Definition of High Risk

The following two groups of individuals are considered as **HIGH RISK** in this guideline.

1. People with the following medical problems or conditions are at **highest** risk of complications.

- Age: The risk for severe illness from COVID-19 increases with age, with older adults at highest risk.
- Cancer
- Chronic kidney disease
- Heart disease, such as heart failure, coronary artery disease
- Obesity with a body mass index (BMI) of 30 or higher
- Sickle cell disease
- Smoking
- Solid organ transplant
- Type 2 diabetes

2. According to the CDC, people with the following medical conditions might be at **higher** risk of severe illness from COVID-19 (or from influenza).

- Asthma (moderate-to-severe)
- Cerebrovascular disease (affects blood vessels and blood supply to the brain)
- Cystic fibrosis
- Hypertension or high blood pressure
- Immunocompromised state (weak immune system) from blood or bone marrow transplant, immune deficiencies, HIV, use of corticosteroids, or use of other immune weakening medicines
- Liver disease
- Neurologic conditions, such as dementia
- Pregnancy
- Pulmonary fibrosis (having damaged or scarred lung tissues)
- Thalassemia (a type of blood disorder)
- Type 1 diabetes

The adult death rate for COVID-19 is approximately 1% to 3%. The death rate is lower in children and younger adults. It is higher in older adults.

### Treatment

Treatment is supportive. Oxygen and IV fluids are used for hospitalized patients. Steroid medications are used in some hospitalized patients.

On May 1, 2020 the U.S. Food and Drug Administration (FDA) granted temporary emergency use authorization (EUA) for the investigational antiviral remdesivir to treat COVID-19. Doctors can use this drug to treat patients who are hospitalized with COVID-19 who are severely ill. More research is needed to determine how well this drug works and for which patients.

None of the other drugs mentioned recently in the news (such as chloroquine) have any proven efficacy in treating or preventing this viral disease.

### Prevention

Social distancing and wearing masks have been proven to help prevent COVID-19.

### **Vaccination**

There currently is no approved vaccine to prevent COVID-19. There are vaccines in development. A COVID-19 vaccine may be available for adults by early 2021.

### **Going Out in Public**

The risk of COVID-19 spread increases with how closely a person interacts with others, how many people there are, and the longer the interaction.

A person should take the following whenever they go out:

- A cloth face-covering
- Hand sanitizer with at least 60% alcohol

The CDC provides additional instructions and information at: <https://www.cdc.gov/coronavirus/2019-ncov/daily-life-coping/going-out.html>.

### **Travel**

- Avoid all non-essential travel.
- If you must travel, go to the CDC website for updates on travel advisories: <https://www.cdc.gov/coronavirus/2019-ncov/travelers>.

### **Ibuprofen and Other NSAID Use for COVID-19**

Many callers have expressed concerns that ibuprofen (or other NSAID) use to treat COVID-19 symptoms may worsen the disease. These concerns originated from a few physicians' comments and have since spread over social media.

To date, there is no scientific evidence (clinical trials or studies) that show that using ibuprofen negatively impacts outcome in COVID-19 patients. We will continue to review any new literature as it is published. The CDC, WHO, AAP and our Infectious Disease expert reviewers continue to approve the use of ibuprofen for COVID-19.

For these reasons, Schmitt-Thompson Clinical Content (STCC) guidelines continue to recommend ibuprofen as an acceptable way to treat high fevers and pain.

- Remind callers that fevers may be beneficial, help fight the infection, and speed recovery.
- Low-grade fevers should not be treated.

If callers remain concerned, they can use acetaminophen for symptoms that warrant treatment.

*Caution:* For suspected COVID-19 patients on oral steroids, such as prednisone, the triager should involve the HCP for a decision about whether the drug can be continued.

### **Other Coronaviruses in Humans**

*Common coronaviruses* can cause colds and upper respiratory symptoms. These can be identified in currently available commercial respiratory testing panels (human coronaviruses HKU1, OC43, 229E, and OC43). These coronaviruses are completely different than the novel coronavirus addressed in this guideline.

Two other coronaviruses that previously have caused serious outbreaks are:

- *MERS-CoV*: Middle East Respiratory Syndrome (MERS)
- *SARS-CoV*: Severe Acute Respiratory Syndrome (SARS)

### Animals and COVID-19

- The main way COVID-19 spreads is from person to person. There is low risk of getting COVID-19 from a pet or other animal.
- It is possible for animals to catch COVID-19 from people. A few pets have tested positive for COVID-19 (including cats and dogs).
- The CDC recommends treating pets like other family members when trying to avoid spreading COVID-19. Do not let pets have close contact with other people or animals outside your household. A sick person should self-isolate and avoid contact with both people and pets.
- Call your vet if your pet gets sick or you have other questions.
- The CDC has more information on COVID-19 and animals at: <https://www.cdc.gov/coronavirus/2019-ncov/daily-life-coping/animals.html>.

### Internet Resources

- *Centers for Disease Control and Prevention (CDC)*: Coronavirus. <https://www.cdc.gov/coronavirus/>.
- *National Institutes of Health (NIH)*: Treatment Guidelines. <https://www.covid19treatmentguidelines.nih.gov/>.
- *Public Health Agency of Canada*: <https://www.canada.ca/en/public-health/services/diseases/coronavirus-disease-covid-19.html>.
- *World Health Organization (WHO)*: Coronavirus. <https://www.who.int/health-topics/coronavirus>.

*Expert Reviewer*: Lisa M. Koonin DrPH, MN, MPH. Senior Advisor in support of the CDC Coronavirus Disease (COVID-19) Response. Centers for Disease Control and Prevention (CDC).

### REFERENCES

1. Abuelgasim H, Albury C, Lee J. Effectiveness of honey for symptomatic relief in upper respiratory tract infections: a systematic review and meta-analysis *BMJ Evid Based Med*. 2020;bmjebm-2020-111336.
2. ACOG Committee. ACOG Committee Opinion No. 753: Assessment and Treatment of Pregnant Women With Suspected or Confirmed Influenza. *Obstet Gynecol*. 2018 Oct;132(4):e169-e173.
3. Blagev DP, Harris D, Dunn AC, Guidry DW, Grissom CK, Lanspa MJ. Clinical presentation, treatment, and short-term outcomes of lung injury associated with e-cigarettes or vaping: a prospective observational cohort study. *Lancet*. 2019 Dec 7;394(10214):2073-2083.
4. Bonow RO, Fonarow GC, O’Gara PT, Yancy CW. Association of Coronavirus Disease 2019 (COVID-19) With Myocardial Injury and Mortality. *JAMA Cardiol*. Published online March 27, 2020.
5. Bonow RO, Fonarow GC, O’Gara PT, Yancy CW. Association of Coronavirus Disease 2019 (COVID-19) With Myocardial Injury and Mortality. *JAMA Cardiol*. Published online March 27, 2020. doi:10.1001/jamacardio.2020.1105
6. Centers for Disease Control and Prevention (CDC). Severe Outcomes Among Patients with Coronavirus Disease 2019 (COVID-19) - United States. February 12 - March 16, 2020. *MMWR Morb Mortal Wkly Rep*. ePub: 18 March 2020.

7. Centers for Disease Control and Prevention (CDC). February 12 - March 28, 2020. Preliminary Estimates of the Prevalence of Selected Underlying Health Conditions Among Patients with Coronavirus Disease 2019 - United States. *MMWR Morb Mortal Wkly Rep.* ePub: 31 March 2020.
8. Centers for Disease Control and Prevention (CDC). February 12 - March 28, 2020. Preliminary Estimates of the Prevalence of Selected Underlying Health Conditions Among Patients with Coronavirus Disease 2019 - United States. *MMWR Morb Mortal Wkly Rep.* ePub: 31 March 2020.
9. Cortegiani A, Ingoglia G, Ippolito M, Giarratano A, Einav S. A systematic review on the efficacy and safety of chloroquine for the treatment of COVID-19. *J Crit Care.* 2020 Mar 10. pii: S0883-9441(20)30390-7. doi: 10.1016/j.jcrc.2020.03.005. [Epub ahead of print].
10. Davey K, Saul T, Russel G, Wassermann J, Quaas J. Application of the Canadian Computed Tomography Head Rule to Patients With Minimal Head Injury. *Ann Emerg Med.* 2018 Oct;72(4):342-350.
11. Deeks JJ, Dinnes J, Takwoingi Y, et.al. Cochrane COVID-19 Diagnostic Test Accuracy Group. Antibody tests for identification of current and past infection with SARS-CoV-2. *Cochrane Database Syst Rev.* 2020 Jun 25;6(6):CD013652. Cochrane COVID-19 Diagnostic Test Accuracy Group.
12. Deutsch A, Blasiak R, Keyes A, Wu J, Marmon S, Asrani F, Moy J, Russo M, McLellan BN. COVID toes: Phenomenon or epiphenomenon? *J Am Acad Dermatol.* 2020 Nov;83(5):e347-e348.
13. Dinnes J, Deeks JJ, Adriano A, et.al. Cochrane COVID-19 Diagnostic Test Accuracy Group. Rapid, point-of-care antigen and molecular-based tests for diagnosis of SARS-CoV-2 infection. *Cochrane Database Syst Rev.* 2020 Aug 26;8:CD013705.
14. Ebell MH, Lundgren J, Youngpairoj S. How long does a cough last? Comparing patients' expectations with data from a systematic review of the literature. *Ann Fam Med.* 2013 Jan;11(1):5-13.
15. Eccles R. Understanding the symptoms of the common cold and influenza. *Lancet Infect Dis.* 2005 Nov;5(11):718-25.
16. Gibani MM, Toumazou C, Sohbati M, et.al. Assessing a novel, lab-free, point-of-care test for SARS-CoV-2 (CovidNudge): a diagnostic accuracy study. *Lancet Microbe.* 2020 Sep 17.
17. Godwin SA, Cherkas DS, Panagos PD, Shih RD, Byyny R, Wolf SJ. Clinical Policy: Critical Issues in the Evaluation and Management of Adult Patients Presenting to the Emergency Department With Acute Headache. *Ann Emerg Med.* 2019 Oct;74(4):e41-e74.
18. Grohskopf LA, Alyanak E, Broder KR, et al. Prevention and Control of Seasonal Influenza with Vaccines: Recommendations of the Advisory Committee on Immunization Practices - United States, 2020-21 Influenza Season. *MMWR Recomm Rep.* 2020;69(8):1-24. Published 2020 Aug 21.
19. Guan WJ, Ni ZY, Hu Y, et al. Characteristics of Coronavirus Disease 2019 in China. *N Engl J Med.* 2020;382(18):1708-1720.
20. Huang C, Wang Y, et.al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* published online. January 24, 2020. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5).
21. Hui DS. Epidemic and Emerging Coronaviruses (Severe Acute Respiratory Syndrome and Middle East Respiratory Syndrome). *Clin Chest Med.* 2017 Mar;38(1):71-86.



22. Jiang F, Deng L, Zhang L, Cai Y, Cheung CW, Xia Z. Review of the Clinical Characteristics of Coronavirus Disease 2019 (COVID-19). *J Gen Intern Med.* 2020;35(5):1545-1549.
23. Lauer SA, Grantz KH, Bi Q, et al. The Incubation Period of Coronavirus Disease 2019 (COVID-19) From Publicly Reported Confirmed Cases: Estimation and Application. *Ann Intern Med.* 2020;172(9):577-582.
24. Lee Y, Min P, Lee S, Kim SW. Prevalence and Duration of Acute Loss of Smell or Taste in COVID-19 Patients. *J Korean Med Sci.* 2020 May 11;35(18):e174.
25. Li Q, Zhou L, Zhou M, Chen Z, et.al. Preliminary Report: Epidemiology of the Avian Influenza A (H7N9) Outbreak in China. *N Engl J Med.* 2013 Apr 24. [Epub ahead of print].
26. Ludvigsson JF. Systematic review of COVID-19 in children show milder cases and a better prognosis than adults. *Acta Paediatr.* 2020 Mar 23. doi: 10.1111/apa.15270. [Epub ahead of print].
27. Mizumoto K, Kagaya K, Zarebski A, Chowell G. Estimating the asymptomatic proportion of coronavirus disease 2019 (COVID-19) cases on board the Diamond Princess cruise ship, Yokohama, Japan, 2020. *Euro Surveill.* 2020;25(10):2000180.
28. Pascarella G, Strumia A, Piliago C, et al. COVID-19 diagnosis and management: a comprehensive review [published online ahead of print, 2020 Apr 29]. *J Intern Med.* 2020;10.1111/joim.13091.
29. Paules CI, Marston HD, Fauci AS. Coronavirus Infections - More Than Just the Common Cold. *JAMA*, Published online January 23, 2020.
30. Pringle JC, et.al. COVID-19 in a Correctional Facility Employee Following Multiple Brief Exposures to Persons with COVID-19 - Vermont, July-August 2020. *MMWR* October 21, 2020 / 69. Early Release.
31. Radonovich LJ Jr, Simberkoff MS3, Bessesen MT, et.al. N95 Respirators vs Medical Masks for Preventing Influenza Among Health Care Personnel: A Randomized Clinical Trial. *JAMA.* 2019 Sep 3;322(9):824-833.
32. Rasmussen SA, Smulian JC, Lednicky JA, Wen TS, Jamieson DJ. Coronavirus Disease 2019 (COVID-19) and Pregnancy: What obstetricians need to know. *Am J Obstet Gynecol.* 2020 Feb 24. pii: S0002-9378(20)30197-6. doi: 10.1016/j.ajog.2020.02.017. [Epub ahead of print].
33. Rothberg MB, Haessler SD, Brown RB. Complications of viral influenza. *Am J Med.* 2008 Apr;121(4):258-64.
34. Rothman RE, Irvin CB, Moran GJ, et.al. Public Health Committee of the American College of Emergency Physicians. Respiratory hygiene in the emergency department. *Ann Emerg Med.* 2006;48(5):570-82.
35. Song Z, Xu Y, et.al. From SARS to MERS, thrusting coronaviruses into the spotlight. *Viruses.* 2019 Jan 14;11(1).
36. Struyf T, Deeks JJ, Dinnes J, Takwoingi Y, et.al. Cochrane COVID-19 Diagnostic Test Accuracy Group. Signs and symptoms to determine if a patient presenting in primary care or hospital outpatient settings has COVID-19 disease. *Cochrane Database Syst Rev.* 2020 Jul 7;7(7):CD013665.
37. Uyeki TM, Bernstein HH, Bradley JS, et.al. Clinical Practice Guidelines by the Infectious Diseases Society of America: 2018 Update on Diagnosis, Treatment, etc. of Seasonal Influenza. *Clin Infect Dis.* 2019 Mar 5;68(6):895-902.

38. Vilke G, Chan T, Bozeman WP, Childers R. Emergency Department Evaluation After Conducted Energy Weapon Use: Review of the Literature for the Clinician. *J Emerg Med.* 2019 Nov;57(5):740-746.
39. Wang AZ, Schaffer JT, Holt DB, Morgan KL, Hunter BR. Troponin Testing and Coronary Syndrome in Geriatric Patients With Nonspecific Complaints: Are We Overtesting? *Acad Emerg Med.* 2020 Jan;27(1):6-14.
40. Wang H, Feng Z, Shu Y, Yu H, et.al. Probable limited person-to-person transmission of highly pathogenic avian influenza A (H5N1) virus in China. *Lancet.* 2008 Apr 26;371(9622):1427-34
41. Wang W, Xu Y, Gao R, Lu R, Han K, Wu G, Tan W. Detection of SARS-CoV-2 in Different Types of Clinical Specimens. *JAMA.* 2020 Mar 11. doi: 10.1001/jama.2020.3786. [Epub ahead of print].
42. Yu IT, Li Y, Wong TW, et al. Evidence of airborne transmission of the severe acute respiratory syndrome virus. *N Engl J Med.* 2004;350(17):1731-1739.

## SEARCH WORDS

2019-NCOV  
ACHES  
BREATHING  
BREATHING DIFFICULTY  
CHILLS  
CORONAVIRUS  
COUGH  
COVID19  
COVID-19  
DIFFICULT BREATHING  
DIFFICULTY BREATHING  
FOREIGN TRAVEL  
LOSS OF SMELL  
LOSS OF TASTE  
NCOV  
NOVEL CORONAVIRUS  
SOB  
TRAVEL  
TROUBLE BREATHING

## AUTHOR AND COPYRIGHT

**Author:** David A. Thompson, MD, FACEP  
**Copyright:** 2000-2020, LaGrange Medical Software, Inc. All rights reserved.  
**Company:** Schmitt-Thompson Clinical Content  
**Content Set:** After Hours Telephone Triage Protocols - Standard | Adult  
**Version Year:** 2020  
**Last Revised:** 11/3/2020  
**Last Reviewed:** 11/3/2020