

DEFINITION

Child has symptoms of COVID-19 (cough, fever, shortness of breath or others) AND:

- **Positive lab test confirms the diagnosis** OR
- **Doctor (or NP or PA) makes a clinical diagnosis** (suspected diagnosis) OR
- **Parent or patient makes suspected diagnosis** based on symptoms consistent with COVID-19 AND any possible close contact with COVID-19 patient within last 2 weeks OR
- **Triage Nurse makes suspected diagnosis** using nurse judgment based on symptoms consistent with COVID-19 AND any possible close contact with COVID-19 patient within last 2 weeks.
Perspective: triage nurses are more qualified to suspect a clinical diagnosis of this infection than parents.
- Confirmation of Diagnosis: COVID-19 testing is now widely available. It should be performed on all the above suspected cases. Testing can confirm which of the suspected cases have a COVID-19 diagnosis.
- **Also Included:** Suspected Influenza calls during the 2021-2022 flu season when flu is also present in the community.
- **Updated:** November 15, 2021 (version 14)

COVID-19 Fully Vaccinated Symptomatic Patients who later Develop COVID-19 Compatible Symptoms:

- COVID-19 vaccines approved by the FDA are highly effective. Research data has confirmed that protective antibody levels are still high at 9 months in most people after completing the vaccine series.
- However, some may develop a mild breakthrough infection.
- This guideline continues to SMAG *fully vaccinated* patients with COVID-19 compatible symptoms and no known or possible exposure to other symptom-based guidelines. This is a practical decision.
- Each call center medical director will need to consider this decision based on changes in variants and vaccines.
- Note: A **fully vaccinated** patient means 2 weeks have passed since the final dose. A **partially vaccinated** patient means incomplete vaccine series or less than 2 weeks have passed since final dose.
- The CDC recommendation for **fully vaccinated asymptomatic** people who have come into close contact with someone with suspected or confirmed COVID-19 is to be tested 5-7 days after exposure and to wear a mask in public indoor settings for 14 days or until they receive a negative test result. No home quarantine is needed.

COVID-19 Main Symptoms (CDC)

COVID-19 should be suspected in people who have 1 or more of the following symptoms (CDC) and have not been fully vaccinated against COVID-19:

- Cough
- Shortness of breath (difficulty breathing)
- Fever or chills
- Loss of smell or taste
- Muscle or body aches
- Headache
- Sore throat

- Runny nose (not from allergies)
- The CDC also includes the following less common symptoms: fatigue (tiredness), nausea, vomiting and diarrhea. In isolation, these symptoms are not very helpful for recognizing COVID-19. Unless there is associated close contact with a COVID-19 patient, these symptoms can usually be triaged and managed in those specific guidelines. So can an isolated headache. For reasons of safety, all respiratory symptoms (such as runny nose and sore throat) are considered COVID-19 until disproven by testing.

Influenza Calls: Preventing the Need to Use 2 Guidelines

Here are the reasons why this guideline can be used simultaneously for calls about patients with suspected COVID-19 and also for those with suspected Influenza:

- **Symptoms** are nearly identical. Cannot differentiate based on symptoms. Only exception: loss of taste or smell is highly specific for COVID.
- **Triage for serious symptoms** or complications is the same. The nurse can triage both at same time.
- **Viral Testing** is the only way to reach an accurate diagnosis. Tests for both are available.
- **Care Advice** is the same. Treat symptoms and stay well hydrated.
- **Oral Antivirals** are available for patients with influenza who also are High Risk for complications.
- **High-Risk patients for Complications:** The long-established list for influenza is similar to the evolving list for patients with COVID-19. It can be used for both.
- **Isolation:** Home isolation is required for 10 days or longer for COVID-19. Isolation for flu is only recommended until the fever is gone for 24 hours or longer. Reason: COVID-19 is far more dangerous than flu.
- **Why COVID-19 Guideline was Chosen to Cover Both:** Influenza is seasonal. COVID-19 is not seasonal. It will not go away in 6 months like influenza.

Preventing the Need to Use 2 Symptom Guidelines to Manage COVID-19 Calls

- **Concern:** Call centers have reported that triage nurses often needed to use 2 guidelines to provide complete triage and care advice for suspected COVID-19 cases. Reason: Many patients have multiple symptoms.
- **Solution:** Triage and targeted care advice have been expanded.
- This guideline now covers telephone care for Cough, Fever, Chills or shaking, Sore throat, Muscle pains, Headaches and Loss of smell or taste.
- Triage nurses should use clinical judgment and consider use of an additional guideline for any important symptoms not addressed in this guideline. Examples are Asthma or Vomiting.

INITIAL ASSESSMENT QUESTIONS

Note to Triage - Respiratory Distress: Always rule out respiratory distress (also known as working hard to breathe or shortness of breath). Listen for grunting, stridor, wheezing, tachypnea in these calls. How to assess: Listen to the child's breathing early in your assessment. Reason: What you hear is often more valid than the caller's answers to your triage questions.

1. COVID-19 DIAGNOSIS: "Who made your COVID-19 diagnosis? Was it confirmed by a positive lab test?"
2. COVID-19 EXPOSURE: "Was there any known exposure to COVID-19 before the symptoms began?" Household exposure or close contact with positive COVID-19 patient outside the home (child care, school, work, play or sports). CDC Definition of close contact: within 6 feet (2 meters) for a total of 15 minutes or more over a 24-hour period.
3. ONSET: "When did the COVID-19 symptoms start?"
4. WORST SYMPTOM: "What is your child's worst symptom?"
5. COUGH: "Does your child have a cough?" If so, ask, "How bad is the cough?"
6. RESPIRATORY DISTRESS: "Describe your child's breathing. What does it sound like?" (e.g.,

- wheezing, stridor, grunting, weak cry, unable to speak, retractions, rapid rate, cyanosis)
7. BETTER-SAME-WORSE: "Is your child getting better, staying the same or getting worse compared to yesterday?" If getting worse, ask, "In what way?"
8. FEVER: "Does your child have a fever?" If so, ask: "What is it, how was it measured, and how long has it been present?"
9. OTHER SYMPTOMS: "Does your child have any other symptoms?" (e.g., chills or shaking, sore throat, muscle pains, headache, loss of smell)
10. CHILD'S APPEARANCE: "How sick is your child acting?" " What is he doing right now?" If asleep, ask: "How was he acting before he went to sleep?"
11. HIGHER RISK for COMPLICATIONS with FLU or COVID-19 : "Does your child have any chronic medical problems?" (e.g., heart or lung disease, diabetes, asthma, cancer, weak immune system, etc. See that List in Background Information. Reason: may need antiviral if has positive test for influenza.)

- Author's note: IAQ's are intended for training purposes and not meant to be required on every call.

TRIAGE ASSESSMENT QUESTIONS

Call EMS 911 Now

Severe difficulty breathing (struggling for each breath, unable to speak or cry, making grunting noises with each breath, severe retractions) (Triage tip: Listen to the child's breathing.)

CA: 50, 27, 28, 25, 1

Slow, shallow, weak breathing

R/O: *respiratory depression with impending apnea*

CA: 50, 27, 28, 25, 1

[1] Bluish (or gray) lips or face now AND [2] persists when not coughing

R/O: *cyanosis and need for oxygen*

CA: 50, 27, 28, 25, 1

Difficult to awaken or not alert when awake (confusion)

R/O: *encephalitis*

CA: 50, 27, 28, 25, 1

Very weak (doesn't move or make eye contact)

R/O: *sepsis or shock*

CA: 50, 27, 28, 25, 1

Sounds like a life-threatening emergency to the triager

CA: 50, 27, 28, 25, 1

See More Appropriate Guideline

Runny nose from nasal allergies

Go to Guideline: *Nasal Allergies (Hay Fever) (Pediatric)*

[1] COVID-19 compatible symptoms BUT [2] NO possible COVID-19 close contact within last 2 weeks for the child (e.g., only child kept at home with vaccinated caregivers)

Go to the specific symptom-based guideline. Reason: COVID-19 unlikely.

[1] Headache is isolated symptom (no fever) AND [2] no known COVID-19 close contact

Go to Guideline: Headache (Pediatric)

[1] Vomiting is isolated symptom (no fever) AND [2] no known COVID-19 close contact

Go to Guideline: Vomiting without Diarrhea (Pediatric)

[1] Diarrhea is isolated symptom (no fever) AND [2] no known COVID-19 close contact

Go to Guideline: Diarrhea (Pediatric)

[1] COVID-19 exposure AND [2] NO symptoms

Go to Guideline: COVID-19 - Exposure (Pediatric)

[1] COVID-19 vaccine series completed (fully vaccinated) AND [2] new-onset of possible COVID-19 symptoms BUT [3] no possible exposure

Go to the specific symptom-based guideline. Reason: COVID-19 unlikely in fully vaccinated person with no possible new exposure to COVID-19.

[1] Had lab test confirmed COVID-19 infection within last 3 months AND [2] new-onset of possible COVID-19 symptoms BUT [3] no possible exposure

Go to the specific symptom-based guideline. Reason: COVID-19 unlikely in previously infected person with no possible new exposure to COVID-19.

[1] COVID-19 vaccine general reaction (fever, headache, muscle aches, fatigue) AND [2] starts within 48 hours of shot (Note: vaccine does not cause respiratory symptoms. Stay here for those symptoms.)

Go to Guideline: Immunization Reactions (Pediatric)

COVID-19 vaccine, questions about

Go to Guideline: Immunization Reactions (Pediatric)

[1] Diagnosed with influenza within the last 2 weeks by a HCP AND [2] follow-up call

Go to Guideline: Influenza (Flu) Follow-up Call (Pediatric)

[1] Household exposure to known influenza (flu test positive) AND [2] child with influenza-like symptoms

Go to Guideline: Influenza (Flu) - Seasonal (Pediatric)

Go to ED Now

[1] Difficulty breathing confirmed by triager BUT [2] not severe (Triage tip: Listen to the child's breathing.)

R/O: pneumonia

CA: 51, 37, 23, 25, 26, 1

Ribs are pulling in with each breath (retractions)

R/O: pneumonia

CA: 51, 37, 23, 25, 26, 1

[1] Age < 12 weeks AND [2] fever 100.4 F (38.0 C) or higher rectally

R/O: sepsis

CA: 51, 37, 23, 25, 26, 33, 1

SEVERE chest pain or pressure (excruciating)

R/O: pneumonia, pleurisy, pulmonary emboli

CA: 51, 37, 23, 25, 26, 31, 1

Go to ED Now (or PCP triage)

[1] Stridor (harsh sound with breathing in) AND [2] present now OR has occurred 2 or more times

CA: 52, 37, 23, 25, 26, 83, 4, 84, 1

Rapid breathing (Breaths/min > 60 if < 2 mo; > 50 if 2-12 mo; > 40 if 1-5 years; > 30 if 6-11 years; > 20 if > 12 years)

R/O: respiratory distress. (Caution: Do not attribute abnormal RR to fever)

CA: 52, 37, 23, 25, 26, 4, 84, 1

[1] MODERATE chest pain or pressure (by caller's report) AND [2] can't take a deep breath

R/O: pneumonia, pleurisy

CA: 52, 37, 23, 25, 26, 4, 31, 84, 1

[1] Fever AND [2] > 105 F (40.6 C) by any route OR axillary > 104 F (40 C)

R/O: serious bacterial infection

CA: 52, 37, 23, 25, 26, 4, 84, 1

[1] Shaking chills (shivering) AND [2] present constantly > 30 minutes

R/O: sepsis

CA: 52, 37, 23, 25, 26, 4, 84, 1

[1] Sore throat AND [2] complication suspected (refuses to drink, can't swallow fluids, new-onset drooling, can't move neck normally or other serious symptom)

CA: 52, 37, 23, 25, 26, 4, 84, 1

[1] Muscle or body pains AND [2] complication suspected (can't stand, can't walk, can barely walk, can't move arm or hand normally or other serious symptom)

CA: 52, 37, 23, 25, 26, 4, 84, 1

[1] Headache AND [2] complication suspected (stiff neck, incapacitated by pain, worst headache ever, confused, weakness or other serious symptom)

CA: 52, 37, 23, 25, 26, 4, 84, 1

[1] Dehydration suspected AND [2] age < 1 year (signs: no urine > 8 hours AND very dry mouth, no tears, ill-appearing, etc.)

CA: 52, 37, 23, 25, 26, 29, 4, 84, 1

[1] Dehydration suspected AND [2] age > 1 year (signs: no urine > 12 hours AND very dry mouth, no tears, ill-appearing, etc.)

CA: 52, 37, 23, 25, 26, 29, 4, 84, 6, 1

Child sounds very sick or weak to the triager

Reason: severe acute illness or serious complication suspected

CA: 52, 37, 23, 25, 26, 84, 1

See HCP within 4 Hours (or PCP Triage)

[1] Wheezing confirmed by triager AND [2] no trouble breathing (Exception: known asthmatic)

Note to triager: Asthmatic children will also need triaging with the Asthma guideline.

CA: 53, 37, 29, 4, 84, 6, 1

[1] Lips or face have turned bluish BUT [2] only during coughing fits

R/O: need for oxygen

CA: 53, 37, 29, 4, 84, 6, 1

[1] Age < 3 months AND [2] lots of coughing

R/O: pneumonia

CA: 53, 37, 23, 25, 26, 84, 6, 1

[1] Crying continuously AND [2] cannot be comforted AND [3] present > 2 hours

R/O: severe otitis or sinusitis

CA: 53, 37, 23, 25, 26, 29, 4, 31, 84, 6, 1

Call PCP Now

[1] SEVERE RISK patient (e.g., immuno-compromised, serious lung disease, on oxygen, heart disease, bedridden, etc) AND [2] suspected COVID-19 with mild symptoms (Exception: Already seen by PCP and no new or worsening symptoms.)

Reason: special chronic diseases at risk for severe pneumonia or sepsis

CA: 59, 4, 7, 6, 1, 49

[1] Age less than 12 weeks AND [2] suspected COVID-19 with mild symptoms

Reason: PCP will decide on needed follow-up care

CA: 59, 33, 7, 6, 1

Multisystem Inflammatory Syndrome (MIS-C) suspected (Fever AND 2 or more of the following: widespread red rash, red eyes, red lips, red palms/soles, swollen hands/feet, abdominal pain, vomiting, diarrhea)

Note: very rare complication

CA: 59, 4, 6, 1

See PCP within 24 Hours

[1] Stridor (harsh sound with breathing in) occurred BUT [2] not present now

Reason: may need Decadron

CA: 54, 37, 29, 5, 30, 4, 89, 24, 7, 84, 38, 1

[1] Continuous coughing keeps from playing or sleeping AND [2] no improvement using cough treatment per guideline

CA: 54, 37, 29, 5, 30, 4, 7, 84, 6, 1, 49

Earache or ear discharge also present

R/O: otitis media

CA: 54, 37, 4, 32, 29, 5, 30, 7, 84, 6, 1, 49

Strep throat infection suspected by triager

Reason: may need Strep test

CA: 54, 37, 4, 42, 43, 7, 84, 6, 1

[1] Age 3-6 months AND [2] fever present > 24 hours AND [3] without other symptoms (no cold, cough, diarrhea, etc.)

R/O: UTI, bacteremia

CA: 54, 37, 4, 7, 84, 6, 1

[1] Age 6 - 24 months AND [2] fever present > 24 hours AND [3] without other symptoms (no cold, diarrhea, etc.) AND [4] fever > 102 F (39 C) by any route OR axillary > 101 F (38.3 C)

R/O: UTI, bacteremia

CA: 54, 37, 4, 7, 84, 6, 1

[1] Fever returns after gone for over 24 hours AND [2] symptoms worse or not improved

R/O: otitis media or sinusitis

CA: 54, 37, 4, 29, 5, 30, 7, 84, 6, 1, 49

Fever present > 3 days (72 hours)

R/O: bacterial superinfection - usually otitis media

CA: 54, 37, 4, 29, 5, 30, 7, 84, 6, 1, 49

[1] Age > 5 years AND [2] sinus pain around cheekbone or eye (not just congestion) AND [3] fever

R/O: sinusitis

CA: 54, 37, 4, 29, 5, 30, 7, 84, 6, 1, 49

Call PCP within 24 Hours

[1] Influenza also widespread in the community AND [2] mild flu-like symptoms WITH FEVER AND [3] HIGH-RISK patient for complications with Flu (See that CDC List)

Reason: may need testing for influenza and COVID-19. If positive for flu, PCP will decide if antiviral meds would be helpful for this patient.

CA: 60, 85, 86, 87, 4, 5, 29, 42, 43, 44, 11, 88, 82, 95, 6, 1, 49

Call PCP when Office is Open

[1] COVID-19 rapid test result was negative BUT [2] caller worried that child has COVID-19 infection AND [3] mild symptoms (cough, fever, or others) continue

Reason: PCP will decide if PCR test is indicated

CA: 61, 93, 3, 7, 11, 39, 12, 4, 40, 5, 41, 29, 90, 30, 42, 43, 44, 45, 95, 6, 1

[1] COVID-19 infection suspected by caller or triager AND [2] mild symptoms (cough, fever, or others) AND [3] no complications or SOB

Reason: arrange COVID-19 testing in PCP office or other site

CA: 61, 13, 3, 7, 12, 11, 39, 4, 40, 5, 41, 29, 90, 30, 42, 43, 44, 45, 46, 95, 19, 6, 1, 49

Home Care

[1] COVID-19 diagnosed by positive lab test AND [2] NO symptoms

CA: 58, 92, 91, 39, 95, 6, 1

[1] COVID-19 diagnosed by positive lab test AND [2] mild symptoms (cough, fever or others) AND [3] no complications or SOB

CA: 58, 80, 3, 7, 39, 4, 40, 5, 41, 29, 90, 30, 42, 43, 44, 45, 46, 82, 11, 95, 6, 1, 49

[1] COVID-19 diagnosed by a doctor (or NP/PA) without lab test AND [2] mild symptoms (cough, fever or others) AND [3] no complications or SOB

CA: 58, 2, 3, 7, 39, 4, 40, 5, 41, 29, 90, 30, 42, 43, 44, 45, 46, 82, 11, 95, 6, 49, 1

COVID-19 Home Isolation, questions about

CA: 58, 7, 39, 11, 8, 95, 6, 1

COVID-19 Prevention, questions about

CA: 58, 8, 47, 48, 9, 10, 11, 82, 95, 6, 1

COVID-19 Testing, questions about

CA: 58, 12, 93, 94, 6, 1

COVID-19 Maternal Illness and Breastfeeding

CA: 58, 14, 35, 7, 39, 15, 1

COVID-19 Disease, questions about

Multisystem Inflammatory Syndrome (MIS-C), questions about

CARE ADVICE (CA) -

1. **Care Advice** given per COVID-19 - Diagnosed or Suspected (Pediatric) guideline.
2. **Reassurance and Education - Doctor (or NP/PA) Suspects COVID-19 and Symptoms are Mild:**
 - Your doctor told you that your child probably has COVID-19 based on the symptoms and close contact.
 - Your child did not receive a lab test for COVID-19. Reason: household contact or other caregiver had positive lab test.
 - From what you have told me, your child's symptoms are mild. They stay that way for most children.
 - What to Expect: Mild symptoms usually last less than 2 weeks. Complications are rare in children. Re-infections are very rare.
 - You don't need to see your doctor unless your child develops trouble breathing or becomes worse in any other way.
 - Here's some care advice that should help.
3. **Treatment of Symptoms:**
 - The treatment is the same whether you have COVID-19, influenza or some other respiratory virus.
 - The only difference for COVID-19 is you need to stay on home isolation until you recover (a minimum of 10 days). Reason: You want to protect other people from getting it. For influenza-like illness, the CDC recommends that people remain at home (isolate) until at least 24 hours after they are free of fever.
 - Treat the symptoms that are bothering you the most.
 - **Note to Triager:** Care Advice is available for Cough, Fever, Chills and Shivering, Runny nose, Sore throat, Muscle pains, Headache and Loss of smell. Only discuss treatment for the caller's main symptoms.
 - There is no anti-viral medication for treating COVID-19 at home. New antiviral treatments have been developed for patients who need to be hospitalized.
 - Antibiotics are not helpful for viral infections.
4. **Fever Medicine and Treatment:**
 - For fever above 102 F (39 C), you may use acetaminophen OR ibuprofen (See Dosage table). Avoid aspirin.
 - For fevers 100-102 F (37.8 to 39 C), fever medicines are not needed. Reason: Fever turns on your body's immune system. Fever helps fight the infection.
 - Exception: If your child also has definite pain, treat it.
 - **Fluids.** Encourage cool fluids in unlimited amounts. Reason: prevent dehydration. Age younger than 6 months, only give formula or breastmilk.
 - **Clothing.** For all children, dress in 1 layer of clothing, unless shivering. For shivering, use a blanket until it stops.
 - **Note to triager about ibuprofen concerns:** Discuss only if caller brings up concerns about ibuprofen. Response: The CDC, WHO, AAP and other experts continue to support the use of ibuprofen (if needed) for patients with COVID-19. They found no scientific evidence to support the claim that ibuprofen made this disease worse.

5. **Homemade Cough Medicine:**

- **Age:** 3 Months to 1 year:
- Give warm clear fluids (e.g., apple juice or lemonade) to thin the mucus and relax the airway. Dosage: 1-3 teaspoons (5-15 ml) four times per day.
- Note to Triage: Option to be discussed only if caller complains that nothing else helps: Give a small amount of corn syrup. Dosage: 1/4 teaspoon (1 ml). Can give up to 4 times a day when coughing. Caution: Avoid honey until 1 year old (Reason: risk for botulism).
- **Age** 1 year and older: Use **Honey** 1/2 to 1 tsp (2 to 5 ml) as needed as a homemade cough medicine. It can thin the secretions and loosen the cough. (If not available, can use corn syrup.) OTC cough syrups containing honey are also available. They are not more effective than plain honey and cost much more per dose.
- **Age** 6 years and older: Use **Cough Drops** (throat drops) to decrease the tickle in the throat. If not available, can use hard candy. Avoid cough drops before 6 years. Reason: risk of choking.
- Don't use OTC cough medicines under 6 years of age. Reason: Cough is a protective reflex.

6. **Call Back If:**

- Shortness of breath occurs
- Difficulty breathing occurs
- Your child becomes worse

7. **Home Isolation Is Needed when You or Your Child are Sick with COVID-19:**
- Isolation means separating sick people with a contagious disease from people who are not sick. (CDC)
 - **Stay Home for Minimum of 10 Days:** Home isolation is needed for at least 10 days after symptoms started.
 - Follow local, state or provincial Department of Health directives.
 - Students should follow their school's COVID-19 policy.
 - Living with a suspected COVID-19 patient implies close contact has occurred.
 - Both patient and any *unvaccinated/partially vaccinated* family members should stay home on isolation and quarantine.
 - **Exceptions for Exposure and NO symptoms:** If you are vaccinated and 2 weeks have passed since your final dose, you do not have to quarantine for 10 days after close contact with a COVID-19 infected person. Also, essential workers who have COVID-19 exposure but do not have any symptoms should talk to your employer. However, anyone exposed to COVID-19 regardless of vaccination status and is asymptomatic should get a COVID-19 test (CDC). See testing care advice for timeframes. Test sooner if symptoms develop.
 - **Ongoing Close Contact (Household):** If living with a person who has a COVID-19 positive test, it implies ongoing exposure. Here is some general guidance: The infected person is contagious for up to 10 days. Living in the same household means all household members continue to be exposed a minimum of 10 days. If a household member develops COVID symptoms, they should get tested within 3 days of onset of symptoms. If a household member does NOT develop symptoms, a test is not needed until 15 to 17 days after the date the first household member's symptoms started or positive test was collected. If a second family member tests positive, the cycle starts over.
 - The patient does not need to be confined to a single room. Reason: Preventing spread of respiratory infections within a home is nearly impossible.
 - The sick person should try to avoid very close contact with other family members. That includes hugging, kissing, sitting next to or sleeping in the same bed. None of this is realistic for young children.
 - Older children and adults with symptoms may consider wearing a mask in common household areas.
 - Note to Triager: Many families have limited options. Triagers should individualize their recommendations for isolation after discussing it with the caller.
 - **Isolation Questions for Your PCP:** Home isolation can be complicated. A parent may need to return to work. Someone in the household may be elderly or have a serious medical problem. If you have additional questions, call your doctor during office hours. Your doctor is the best resource for up-to-date information on COVID-19.

8. **COVID-19 - How to Protect Yourself and Family from Catching It - The Basics:**
 - Get the COVID-19 vaccine. It is your best protection against this serious infection.
 - Avoid close contact with people outside your family unit. Avoid closed spaces (indoors) when possible and all crowds (even outdoors).
 - Always wear a mask when you leave your home. Also, observe social (safe) distancing.
 - **Wash hands often with soap and water (very important).** Always do before you eat.
 - Use an alcohol-based hand sanitizer if water is not available. Remember: soap and water work better.
 - Don't touch your eyes, nose or mouth unless your hands are clean. Germs on the hands can get into your body this way.
 - Don't share glasses, plates or eating utensils.
 - No longer shake hands. Greet others with a smile and a nod.
 - If your child needs to be seen for an urgent medical problem, do not hesitate to go in. ERs and urgent care sites are safe places. They are well equipped to protect you against the virus. For non-urgent conditions, talk to your doctor's office first.
9. **Keep Your Body Strong:**
 - Get your body ready to fight the COVID-19 virus.
 - Get enough sleep (very important)
 - Keep your heart strong. Walk or exercise every day. Take the stairs. Caution: avoid physical exhaustion.
 - Stay well hydrated.
 - Eat healthy meals. Avoid overeating to deal with your fears.
 - Avoid the over-use of anti-fever medicines. Fever fights infections and ramps up your immune system.
10. **Keep Your Mind Positive:**
 - **Live in the present, not the future.** The future is where your needless worries live.
 - **Stay positive.** Use a mantra to reduce your fears, such as "I am strong".
 - **Get outdoors.** Take daily walks. Go to a park if you have one. Being in nature is good for your immune system.
 - **Show love.** As long as they are well, hug your children and partner frequently. Speak to them in a kind and loving voice. Love strengthens your immune system.
 - **Stay in touch.** Use regular phone calls and video chats to stay in touch with those you love.

11. **How to Protect Others - When You or Your Child are Sick with COVID-19:**
- **Stay Home for Minimum of 10 Days:** Home isolation is needed for at least 10 days after symptoms started.
 - Follow local, state or provincial Department of Health directives.
 - Students should follow their school's COVID-19 policy.
 - Do Not go to stores, restaurants, places of worship or other public places. Avoid public transportation or ride sharing. Do Not allow any visitors (such as friends). Leave the house only if you need to seek medical care.
 - **Cover the Cough:** Cough and sneeze into your shirt sleeve or inner elbow. Don't cough into your hand or the air. If available, sneeze into a tissue and throw it into trash can.
 - **Wash Hands often with Soap and Water:** After coughing or sneezing are important times.
 - **Don't Share Personal Household Items:** Don't share glasses, plates or eating utensils.
 - **Wear a Mask:** Wear a face mask when around others or if you have to go to a medical facility.
 - Carefully avoid any contact with the elderly and people with weak immune systems or other chronic health problems.
12. **COVID-19 Diagnostic Testing:**
- Testing is the only way to know for sure that your child has COVID-19. You can't tell by symptoms. Reason: Most respiratory viruses cause similar symptoms.
 - Testing is now widely available without a doctor's order. Exception: age less than 3. Where to get a test can be different for some communities.
 - Some doctors can do COVID-19 tests in their office. Many retail clinics and urgent care centers perform COVID-19 testing. Even pharmacies (such as CVS and Walgreens) now perform drive-thru testing on children age 3 and older. Visit their website to schedule a test. Testing is also available at some local and state public health departments. Self-tests (such as Abbot BinaxNow) for use at home are now available in some drugstores (such as CVS, Walgreens).
 - **Diagnostic Tests:** These are performed on nasal or mouth secretions and tell us if your child has a COVID-19 infection now. The type of diagnostic tests that are available continues to improve.
 - **Tests for COVID-19: Recommended Timing (CDC):**
 - **Symptomatic patients** - get a test immediately (or at least within 3 days of onset of symptoms.)
 - **Asymptomatic Unvaccinated or Partially Vaccinated patients with a COVID-19 close contact** - Get a COVID-19 test immediately (within 24 hours). If the test is negative, the test should be repeated 5 to 7 days after exposure. Test sooner if symptoms develop.
 - **Asymptomatic Fully Vaccinated patients with a COVID-19 close contact** - Get a test on day 5-7 after exposure. Test sooner if symptoms develop.
 - **Negative Tests:** After a negative test, a repeat test is sometimes needed. Reason: A test may be falsely negative; for example, if a person gets the test too soon after exposure. Further, if a person is exposed again or develops symptoms suggestive of COVID-19, then repeat viral testing should be performed. Repeat testing with a PCR test may also be indicated after a negative rapid test.
 - **Repeat Positive Tests:** After a positive test, repeat tests are not recommended. Repeat testing with a PCR test is not indicated after a positive rapid test. Even after it is safe to stop isolation (usually 10 days), tests may stay positive for up to 90 days. A positive test does not mean the patient can spread the infection once the required isolation period is completed.
 - If you have more testing questions, call your doctor during office hours.

13. **Caller Suspects COVID-19 and Symptoms are Mild:**
 - You suspect your child has COVID-19 because they have developed symptoms that match. Cough, sore throat and fever are the most common symptoms. In addition, you think there could have been contact with someone who had an active COVID-19 infection.
 - Getting a COVID-19 lab test is the only way to know for sure. Getting the test is not urgent.
 - From what you have told me, your child's symptoms are mild. They stay that way for most children.
 - Here's some care advice that should help.

14. **Breastfeeding and COVID-19:**
 - Breastfeeding experts recommend you continue to breastfeed even if you are sick with COVID-19. (AAP)
 - Wash your hands before feeding your baby.
 - The CDC recommends wearing a face mask or covering. Be careful to avoid coughing on your baby.
 - Breastmilk gives beneficial antibodies your body is making against this illness to your baby. This should provide some protection against this illness for your baby, like it does for influenza and most other viral illnesses.
 - Research has proven that the virus is not passed through breastmilk.
 - Breastfeeding mothers are also encouraged to get the COVID-19 vaccine. (CDC) After a few weeks, the breastmilk will contain protective antibodies against COVID-19.

15. **Call Back If:**
 - Breastfeeding isn't going well
 - Your baby becomes sick

16. **Trusted Sources for Accurate Information - CDC and AAP:**
 - To meet the extreme demand for COVID-19 information, when possible, find your answers online. Here are the most reliable websites:
 - CDC website: <https://www.cdc.gov/coronavirus>.
 - American Academy of Pediatrics parent website: www.healthychildren.org
 - Nurse advice lines and medical call centers are needed for sick patient calls.

17. **COVID-19 Outbreak:**
 - COVID-19 stands for Coronavirus disease 2019.
 - Cause: The name of the new virus is SARS-CoV-2.
 - An outbreak of this infection began in Wuhan, China in early December 2019.
 - The first COVID-19 patient in the United States was reported on January 21, 2020.
 - The first COVID-19 patient in Canada was reported on January 31, 2020.
 - The World Health Organization (WHO) declared COVID-19 a global pandemic on March 11, 2020.
 - In the summer and fall of 2021, the Delta variant has become the most common COVID-19 variant.
 - The Centers for Disease Control and Prevention (CDC) is considered the source of truth. This continues to be a changing situation and recommendations from the CDC are being updated regularly. See:
 - <https://www.cdc.gov/coronavirus>

18. **COVID-19 Symptoms:**
- COVID-19 coronavirus most often causes a respiratory illness. The most common symptoms are cough and fever. Some patients progress to shortness of breath.
 - Other common symptoms are chills, shivering (shaking), runny nose, sore throat, muscle pain, headache, loss of smell and taste.
 - The CDC also includes the following less common symptoms: fatigue (tiredness), nausea, vomiting and diarrhea.
 - Some people may have minimal symptoms or even have no symptoms (asymptomatic).
19. **COVID-19 - Exposure Risk Factors:**
- Here are the main risk factors for getting sick with COVID-19:
 - Close contact with a person who tested positive for COVID-19 AND contact occurred while they were ill. Close contact is defined as being within 6 feet (2 meters) for a total of 15 minutes or more over a 24-hour period. Prolonged close contact would extend the risk to the 48 hours prior to the person becoming ill with symptoms. This includes living with someone infected with COVID-19.
 - Living in or travel to an area where there is high community spread of COVID-19 also carries some risk.
 - International travel: The CDC (<https://www.cdc.gov/coronavirus>) has the most up-to-date list of where COVID-19 outbreaks are highest.
 - Not being fully vaccinated
20. **COVID-19 - How it is Spread:**
- COVID-19 is spread from person to person.
 - The virus spreads when respiratory droplets produced when a person coughs, sneezes, shouts or sings. The infected droplets can then be inhaled by a nearby person or land on the surface of their face or eyes. Droplets fall quickly to the floor or ground. This is how most COVID is spread.
 - Most infected people also have respiratory secretions on their hands. These secretions get transferred to healthy people on doorknobs, faucet handles etc. The virus then gets transferred to healthy people when they touch their face or rub their eyes. This is a less common cause of spread.
 - These methods are how most respiratory viruses spread.
 - Aerosols are tiny, invisible particles that can float in the air for 1 to 2 hours. They only occur in a closed room with poor ventilation. Aerosols are a rare cause of COVID-19 transmission. (CDC and WHO).
21. **COVID-19 - Travel:**
- Avoid all non-essential air travel.
 - Travel is much safer for people who are vaccinated.
 - The Centers for Disease Control and Prevention (CDC) maintains a website with the latest recommendations regarding travel and your health.
 - Currently the CDC recommends against travel to many geographic areas with widespread and ongoing spread of COVID-19. See current list at <https://wwwnc.cdc.gov/travel/>.

22. **COVID-19 - Other Facts:**

- **Incubation Period:** average 5 days (range 2 to 14 days) after coming in contact with the secretions of a person who has COVID-19.
- **No Symptoms but Infected:** Over 30% of infected adult patients have no symptoms (asymptomatic patients). Children and teens are even more likely to have no symptoms. Such patients do however spread the disease and most develop protective antibodies (immunity).
- **Mild Infections:** 80% of adults with symptoms have a mild illness, much like normal flu or a bad cold. The symptoms usually last 2 weeks.
- **Severe Infections:** 20% of unvaccinated adults with symptoms develop trouble breathing from viral pneumonia. Many of these need to be admitted to the hospital. About 2% of unvaccinated children with COVID-19 need to be admitted to the hospital. About 10% of unvaccinated teens need hospitalization. About 3% require ICU care. (CDC). People with complications generally recover in 3 to 6 weeks. Severe infections are rare in people who are vaccinated.
- **Deaths:** Children generally have a mild illness and recover quickly. Pediatric deaths are very rare. (CDC) Older adults, especially those with chronic lung disease, heart disease, diabetes, obesity or weak immune systems, have the highest death rates. The overall death rate is around 2 per 1000 people. Over 90% of deaths occur in people who are not vaccinated.
- **Vaccine:** Safe and effective vaccines are available. Some vaccines are 2 doses, given 3-4 weeks apart. Others are a single dose. Similar to flu shots, they will probably provide protection for 6 to 9 months. At this time, vaccines have been tested and are FDA approved for 5 years and older. The COVID-19 vaccine will reduce the chance of your child getting COVID-19. The vaccine prevents almost all hospital admissions, ICU care and deaths.
- **"Breakthrough Cases":** These are COVID-19 infections that bypass vaccine protection. They are rare and many are asymptomatic.
- **Treatment:** New treatments for severe COVID-19 are available. They are mainly used on hospitalized patients and are given in a vein (IV). **Caution** - only discuss the following if caller asks about the new anti-viral pill (paxlovid): Paxlovid is given by mouth during the first 3 days of symptoms to prevent serious complications. It awaits FDA approval and will initially be used for adults at high risk for severe disease.
- **Prevention:** The COVID-19 vaccine is the best way to prevent infections. Face masks, social (safe) distancing and extra handwashing are also proven to help prevent disease. **Caution** - only discuss the following if caller asks about monoclonal antibody therapy: Is available for those 12 and older at high risk for severe disease who are 1) asymptomatic and had a recent close contact exposure OR 2) have confirmed COVID-19 mild symptoms. It is usually given IV to prevent progression and complications. People hospitalized with COVID-19 are not eligible.

23. **Note to Triage - Triage Nurse Should Notify Emergency Department (ED):**

- Tell them you are sending a patient with suspected diagnosis of COVID-19 who is getting worse and inform them of patient's symptoms.
- Reason: So ED can make plans to prevent COVID-19 spread to others in the hospital.
- Also determine the best means of transportation.

24. **First Aid For Stridor:**
 - For stridor (harsh sound with breathing in) or constant coughing:
 - Breathe warm mist in a foggy bathroom with the hot shower running for 20 minutes. Other options: a wet washcloth held near the face or a humidifier containing warm water.
 - Caution: avoid very hot water or steam which could cause burns or high body temperatures.
 - If warm mist fails, breathe cool air by standing near an open refrigerator or taking outside for a few minutes if the weather is cold.
 - What to Expect. The stridor should go away with warm mist. The cough and hoarse voice won't.

25. **Cover Your Mouth and Nose - Wear a Mask:**
 - Exception: Less than 2 years or face covering increases difficulty breathing.
 - Cover the patient's mouth and nose loosely with a disposable tissue (e.g., Kleenex or paper towel), a washcloth or bandana.
 - Have patient wear a disposable face mask if you have one.
 - Ask for a face mask on arrival.

26. **Announce COVID-19 Diagnosis on Arrival in ED:**
 - Tell the first hospital worker you meet that your child probably has (or does have) COVID -19.
 - Tell them you were referred because of trouble breathing or other serious symptoms.

27. **Tell the Ambulance Dispatcher about COVID-19 Diagnosis:**
 - When you call 911, tell the dispatcher that your child probably has COVID-19.

28. **Tell Ambulance Medics about COVID-19 Diagnosis:**
 - Tell the paramedic right away that your child may have COVID-19.
 - Paramedics should call ahead to the ED to let them know.

29. **Coughing Fits or Spells - Warm Mist and Fluids:**
 - Breathe warm mist (such as with shower running in a closed bathroom).
 - Give warm clear fluids to drink. Examples are apple juice and lemonade. Don't use warm fluids before 3 months of age.
 - Amount. If 3 - 12 months of age, give 1 ounce (30 ml) each time. Limit to 4 times per day. If over 1 year of age, give as much as needed.
 - Reason: Both relax the airway and loosen up any phlegm.
 - What to Expect: The coughing fit should stop. But, your child will still have a cough.

30. **Humidifier:**
 - If the air is dry, use a humidifier in the bedroom (Reason: dry air makes coughs worse).
 - Avoid menthol vapors (Reason: makes coughs worse).

31. **Pain Medicine:**
 - For pain relief, give acetaminophen every 4 hours **Or** ibuprofen every 6 hours as needed. (See Dosage table.)
 - Note to triager about ibuprofen concerns: Discuss only if caller brings up concerns about ibuprofen. Response: The CDC, WHO and other experts continue to support the use of ibuprofen (if needed) for patients with COVID-19. They found no scientific evidence to support the claim that ibuprofen made this disease worse.

32. **Cold or Hot Pack for Ear Pain:**
 - Apply a cold pack or a cold wet washcloth to outer ear for 20 minutes to reduce pain while medicine takes effect.
 - Note: Some children prefer local heat for 20 minutes.
 - **Caution:** cold or hot pack applied too long could cause frostbite or burn.
33. **Fever Under 3 Months Old - Don't Give Fever Medicine:**
 - Don't give any acetaminophen before being seen.
 - Need accurate documentation of temperature in medical setting to decide if fever is really present. (Reason: may require septic work-up.)
34. **Call Back If:**
 - Your child becomes worse
35. **Wash Your Hands with Soap and Water:**
 - Wash your hands and face frequently with soap and water.
36. **Call Back If:**
 - You have other questions
37. **Note to Triage: If Available, Refer for Re-Triage by a Doctor (or NP/PA):**
 - During this pandemic, the medical community is trying extra hard to prevent unnecessary ED referrals of suspected COVID-19 patients. Parents also prefer this. Re-triage by a physician has been shown to reduce ED referrals. Here are some resources that may be available in your community:
 - **PCP telephone re-triage:** Some PCPs (primary care providers) want to provide re-triage before any of their non-emergent patients are referred to an ED. This requires their approval.
 - **ED telephone re-triage:** Some EDs provide a telephone triage service for patients referred in.
 - **Telemedicine re-triage:** Many practices and some hospitals now offer a telemedicine (virtual visit) service. Telemedicine has the highest rate of providing a definitive diagnosis and care without an in-person visit.
38. **Call Back If:**
 - Stridor occurs again
 - Shortness of breath occurs
 - Difficulty breathing occurs
 - Your child becomes worse

39. **Stopping Home Isolation (CDC):**
- Symptomatic patients must meet 3 criteria: [1] Fever gone for at least 24 hours off fever-reducing medicines AND [2] Cough and other symptoms must be improved AND [3] Symptoms started more than 10 days ago.
 - Asymptomatic patients who don't develop symptoms: must stay at home until 10 days have passed since the date the positive COVID-19 test was done (test specimen was collected).
 - Exposed *unvaccinated/partially vaccinated* person: follow the same rules as above. Period of quarantine starts on the date of last exposure and goes for 10 days.
 - Exception for *fully vaccinated* exposed persons with NO symptoms: If it has been 2 weeks since your final vaccination dose, you do not have to quarantine for 10 days after close contact with a COVID-19 infected person.
 - Shorter quarantine option for asymptomatic people: If they get a negative COVID-19 lab test on day 5 to 7 after exposure, can leave quarantine after day 7. (CDC). This helps essential workers return to the work force.
 - Repeat diagnostic tests: After a positive test, repeat tests are not recommended. Even after it is safe to stop isolation, tests may stay positive for up to 90 days.
 - If unsure it is safe for you to leave isolation, call your PCP.
40. **Chills, Shivering and Rigors - Treatment:**
- Shivering occurs when the body needs to raise its core temperature quickly. Shivering generates body heat until the level of fever that the brain needs to fight the infection is reached.
 - Whether or not you take a fever-reducing medicine, here are some ways to stop the shivering:
 - **Blanket.** Wrap the patient in a warm blanket.
 - **Warm bath.** For severe shivering (rigors), the quickest way to get the fever level up is to take a warm bath or shower. Once the fever peaks, the shivering or rigors will stop.
 - **Fluids.** Drink extra fluids to improve hydration and circulation.
41. **OTC Cough Medicine - DM:**
- OTC cough medicines are not recommended. (Reason: no proven benefit for children.)
 - Honey has been shown to work better. (Caution: Avoid honey until 1 year old.)
 - If the caller insists on using one and the child is over 6 years old, use one with dextromethorphan (DM).
 - Follow the instructions on the package.
 - Indication: Give only for severe coughs that interfere with sleep, school or work.
 - Don't use under 6 years of age. Reason: cough is a protective reflex.
42. **Sore Throat Pain Relief:**
- Here are some tips on treating a sore throat:
 - Age over 1 year: Can sip warm fluids such as chicken broth or apple juice. Some children prefer cold foods such as popsicles or ice cream.
 - Age over 6 years: Can also suck on hard candy or lollipops. Butterscotch seems to help.
 - Age over 8 years: Can also gargle. Use warm water with a little table salt added. A liquid antacid can be added instead of salt. Use Mylanta or the store brand. No prescription is needed.
 - Pain medicine: Use if pain interferes with swallowing. Not needed for mild pain.

43. **Sore Throat - Fluids and Soft Diet:**
- Try to get your child to drink adequate fluids.
 - Goal: Keep your child well hydrated.
 - Cold drinks, milk shakes, popsicles, slushes, and sherbet are good choices.
 - Solid Foods: Offer a soft diet. Also avoid foods that need much chewing. Avoid citrus, salty, or spicy foods.
 - Note: Fluid intake is much more important than eating any solid foods.
44. **Muscle Pains - Treatment:**
- Here are some tips on treating muscle pains and body aches:
 - **Massage:** Gently massage any sore muscles.
 - **Stretching:** Gently stretch any sore muscles.
 - **Apply Heat:** Use a heat pack, heating pad or warm wet washcloth. Do this for 10 minutes 3 times per day.
 - **Warm bath:** For widespread muscle pains, consider a warm bath for 20 minutes 2 times a day. Gently exercise the sore muscles under water.
 - **Pain medicine:** For widespread body aches, give acetaminophen every 4 hours OR ibuprofen every 6 hours as needed. (See Dosage table.) Not needed for mild aches.
45. **Headache - Treatment:**
- Here are some tips for treating a headache:
 - **Pain medicine:** Give acetaminophen every 4 hours OR ibuprofen every 6 hours as needed. (See Dosage table.) Not needed for mild headaches.
 - **Cold pack:** Apply a cold wet washcloth or cold pack to the forehead for 20 minutes.
 - **Massage:** Stretch and massage any tight neck muscles.
46. **Loss of Smell and Taste:**
- Losing the sense of smell and taste can be an early symptom of COVID-19.
 - It is strong evidence for having COVID.
 - In 50% of patients, these senses return within 1 to 3 weeks.
 - In 85%, they return within 6 months.
 - Most of the others recover by 1 year.
 - If symptoms persist, it should not delay the end of isolation.
47. **Social (Safe) Distancing and COVID-19 Prevention:**
- Avoid any contact with people known to have COVID-19 infection. Avoid talking to or sitting close to them.
 - **Social (Safe) Distancing:** Try to stay at least 6 feet (2 meters) away from anyone who is sick, especially if they are coughing. Also called physical distancing. Avoid crowds because you can't tell who might be sick.
 - If COVID-19 is widespread in your community, try to stay 6 feet away from everyone outside your family unit.
 - **Stay at Home Orders:** Follow any stay at home (stay in place) orders in your community. Leave your home only for essential needs such as buying food or seeking medical care.
 - **After Stay at Home Orders are Lifted:** Continue social distancing. Also wear a mask when entering any public building or outdoor crowded area. These precautions will be needed for many months. Your state public health department will decide when they are no longer needed.

48. **Face Masks and COVID-19 Prevention:**
- **Overview:** Face masks are essential for reducing the spread of COVID-19. They will also reduce the spread of influenza. People with COVID-19 can have no symptoms, but still spread the virus.
 - Because of the Delta variant (and other possible future variants) recommendations for wearing masks are pretty much the same for people who are vaccinated or unvaccinated. Mask wearing is even more important if you are in an area of high COVID-19 spread or if you have a weak immune system.
- People Who Are Well (Not Sick With COVID-19) Should Wear Masks If:**
- You are in indoor public spaces (such as a church or a grocery store).
 - You are in a crowded outdoor setting (e.g., concert, music festival, rally).
 - You are traveling on a plane, bus, train, or other form of public transportation or in transportation hubs such as airports and train stations.
 - You must be around someone who has symptoms of COVID-19 or has tested positive for COVID-19.
- People Who Are Sick With COVID-19 Must Wear Masks If:**
- You need to leave the home. Example: for medical visits. Patients with trouble breathing in a mask can consider a loose face covering such as a bandana.
 - You are around other people or animals (such as pets).
- Exceptions to Masks:**
- Face coverings are **NOT** recommended for **children under 2 years**.
 - Face mask or covering is optional if outdoors and you can avoid being within 6 feet (2 meters) of other people. Some examples are an outdoor walk or run.
- How to Select and Use a Face Mask:**
- Make sure your mask fully covers your nose and mouth. It should fit snugly under your chin and against the sides of your face.
 - More information on how to select and use a mask is available at: <https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/diy-cloth-face-coverings.html>.
49. **Asthma - See Additional Guideline:**
- Note to Triager: For patients with COVID-19 or possible influenza who also have a diagnosis of Asthma, ask: "Are you taking asthma meds to manage any wheezing or coughing?"
 - If yes, they need additional triage and care advice. Use the appropriate Asthma guideline.
50. **Call EMS 911 Now:**
- Your child needs immediate medical attention. You need to hang up and call 911 (or an ambulance).
 - Triager Discretion: I'll call you back in a few minutes to be sure you were able to reach them.
51. **Go To ED Now:**
- Your child needs to be seen in the Emergency Department immediately.
 - Go to the ED at _____ Hospital.
 - Leave now. Drive carefully.
52. **Go To ED Now (or PCP Triage):**
- **If No PCP (Primary Care Provider) Second-Level Triage:** Your child needs to be seen within the next hour. Go to the ED/UCC at _____ Hospital. Leave as soon as you can.
 - **If PCP Second-Level Triage Required:** Your child may need to be seen. Your doctor (or NP/PA) will want to talk with you to decide what's best. I'll page the on-call provider now. If you haven't heard from the provider (or me) within 30 minutes, go directly to the ED/UCC at _____ Hospital.

53. **See HCP Within 4 Hours (or PCP triage):**
- **If Office Will Be Open:** Your child needs to be seen within the next 3 or 4 hours. Call your doctor's (or NP/PA) office as soon as it opens.
 - **If Office Will Be Closed and No PCP (Primary Care Provider) Second-Level Triage:** Your child needs to be seen within the next 3 or 4 hours. A nearby Urgent Care Center (UCC) is often a good source of care. Another choice is to go to the ED. Go sooner if your child becomes worse.
 - **If Office Will Be Closed and PCP Second-Level Triage Required:** Your child may need to be seen. Your doctor (or NP/PA) will want to talk with you to decide what's best. I'll page the on-call provider now. If you haven't heard from the provider (or me) within 30 minutes, call again. **Note:** If on-call provider can't be reached, send to UCC or ED.
- Note to Triager:**
- Use nurse judgment to select the most appropriate source of care.
 - Consider both the urgency of the patient's symptoms AND what resources may be needed to evaluate and manage the patient.
- Sources of Care:**
- **ED:** Patients who may need surgery or hospital admission need to be sent to an ED. So do most patients with serious symptoms or complex medical problems.
 - **UCC:** Some UCCs can manage patients who are stable and have less serious symptoms (e.g., minor illnesses and injuries). The triager must know the UCC capabilities before sending a patient there. If unsure, call ahead.
 - **OFFICE:** If patient sounds stable and not seriously ill, consult PCP (or follow your office policy) to see if patient can be seen NOW in office.
54. **See PCP Within 24 Hours:**
- **If Office Will Be Open:** Your child needs to be examined within the next 24 hours. Call your child's doctor (or NP/PA) when the office opens and make an appointment.
 - **If Office Will Be Closed:** Your child needs to be examined within the next 24 hours. A clinic or an urgent care center is often a good source of care if your doctor's office is closed or you can't get an appointment.
 - **If Patient Has No PCP:** Refer patient to a clinic or urgent care center. Also try to help caller find a PCP (medical home) for future care.
- Note to Triager:**
- Use nurse judgment to select the most appropriate source of care.
 - Consider both the urgency of the patient's symptoms AND what resources may be needed to evaluate and manage the patient.
55. **See PCP Within 3 Days:**
- Your child needs to be examined within 2 or 3 days.
 - **PCP Visit:** Call your doctor (or NP/PA) during regular office hours and make an appointment. A clinic or urgent care center are good places to go for care if your doctor's office is closed or you can't get an appointment. **Note:** If office will be open tomorrow, tell caller to call then, not in 3 days.
 - **If Patient Has No PCP (Primary Care Provider):** Try to help caller find a PCP for future care (e.g., use a physician referral line). Having a PCP or "medical home" means better long-term care.

56. **See PCP Within 2 Weeks:**
- Your child needs an evaluation for this ongoing problem within the next 2 weeks.
 - **PCP Visit:** Call your child's doctor (or NP/PA) during regular office hours and make an appointment.
 - **If Patient Has No PCP (Primary Care Provider):** A primary care clinic is where you need to be seen for chronic health problems. **Note:** Try to help caller find a PCP (e.g., use a physician referral line). Having a PCP or 'medical home' means better long-term care.
58. **Home Care:**
- You should be able to treat this at home.
59. **Call PCP Now:**
- You need to discuss this with your child's doctor (or NP/PA).
 - I'll page the on-call provider now. If you haven't heard from the provider (or me) within 30 minutes, call again.
60. **Call PCP Within 24 Hours:**
- You need to discuss this with your child's doctor (or NP/PA) within the next 24 hours.
 - **If Office Will Be Open:** Call the office when it opens tomorrow morning.
 - **If Office Will Be Closed:** I'll page the on-call provider now. Exception: From 9 pm to 9 am. Since this isn't urgent, we'll hold the page until morning.
61. **Call PCP When Office Is Open:**
- You need to discuss this with your child's doctor (or NP/PA) within the next few days.
 - Call the office when it is open.
80. **Lab Test Confirmed COVID-19 AND Symptoms are MILD:**
- Your child had a lab test for COVID-19 and it came back positive.
 - A positive result on a PCR test or rapid self-test kit is highly accurate for diagnosing COVID-19. It is highly likely that your child has COVID-19.
 - From what you have told me, your child's symptoms are mild. They stay that way for most children.
 - You don't need to see your doctor unless your child develops trouble breathing or becomes worse in any other way.
 - Here's some care advice that should help.
81. **Multisystem Inflammatory Syndrome (MIS-C):**
- MIS-C is a very rare complication of COVID-19. In general, COVID-19 continues to be a mild disease in children. It cannot be predicted who will get this complication.
 - Prevention: MIS-C can be prevented by getting your child vaccinated against COVID-19.
 - The most common symptoms are fever, a red rash, abdominal pain with vomiting and diarrhea. Half of the patients develop trouble breathing. Some children become confused or overly sleepy. Always has multiple symptoms.
 - Onset of symptoms: Usually about 4 weeks after a COVID-19 infection and apparent recovery.
 - Peak age: 8 years. Age range: 6 months to 21 years.
 - Treatment: Most patients with MIS-C need to be admitted to the hospital. MIS-C is treatable with medications, including IV immune serum globulin and steroids.
 - Prognosis: Most children with MIS-C have a full recovery. The death rate is about 1 per 100.

82. **Flu and COVID-19 Vaccines:**
- Protect your family from influenza by getting your annual flu vaccine. This year is more important than any.
 - Reason: Getting COVID-19 while you also have or are recovering from the flu may increase the chances of getting severe symptoms.
 - Get the COVID-19 for your child as soon as your child is eligible.
 - Both the flu and the COVID-19 vaccines can be given to children at the same time.
83. **Warm Mist:**
- Before going in to be seen, breathe warm mist in a foggy bathroom for 10 minutes (Note: may also help severe croupy cough).
 - Then inhale warm mist from a wet washcloth while driving in (if this is practical).
84. **How to Protect Others When You or Your Child are Sick:**
- **After You are Seen - Stay Home:** Stay home if you are sick. If you are sick, you are contagious to others. At your medical visit, your provider will discuss home isolation timeframes.
 - **Do NOT go to School or Work:** Do Not go to stores, restaurants, places of worship or other public places. Avoid public transportation or ride sharing.
 - **Avoid Visitors:** Do Not allow any visitors (such as friends). Leave the house only if you need to seek additional medical care.
 - **Wear a Mask:** Always wear a face mask if you have to leave your home. Also try to maintain safe distancing from others (6 feet).
 - **Cover the Cough:** Cough and sneeze into your shirt sleeve or inner elbow. Don't cough into your hand or the air. If available, sneeze into a tissue and throw it into trash can.
 - **Wash Hands Often with Soap and Water:** After coughing or sneezing are important times.
85. **Note to Triager - Contacting the PCP on Weekends and Holidays:**
- On these days, page the on-call PCP during the day up until 5 PM.
 - After 5 PM, page them the next morning after 8 AM.
 - Reason: if antivirals are needed, it is best to start them within 48 hours.
86. **Prescription Antiviral Drugs For Influenza:**
- Antiviral drugs (such as Tamiflu) can be helpful for treating the influenza virus.
 - The benefits are limited: Tamiflu usually reduces the time your child is sick by 1 to 1.5 days. It reduces the symptoms, but does not make them go away.
 - To be helpful, antiviral drugs (such as Tamiflu) should be started within 48 hours of the onset of flu symptoms. If the flu symptoms or fever started more than 48 hours ago, they are not useful. (Exception: children with severe disease).
 - Tamiflu also can have side effects: Vomiting in 10% of children.
 - Most normal children do not need an antiviral drug.
 - The CDC recommends they be used for: [1] any patient with severe symptoms AND [2] for most **High-Risk** children with underlying medical problems (see that list).
 - The CDC doesn't recommend antiviral drugs for **Low-Risk** children with normal influenza. These children recover with supportive care of their symptoms.
 - Also, it is not used to prevent flu. Reason: You would need to take the medicine every day for months.

87. **High-Risk Children For Complications Of The Flu:**
- Lung disease (such as asthma)
 - Heart disease (such as congenital heart disease)
 - Neuromuscular disease (such as muscular dystrophy, epilepsy)
 - Cancer or weak immune system conditions
 - Diabetes, sickle cell disease, kidney disease **Or** other chronic disease
 - Down's syndrome
 - Diseases requiring long-term aspirin therapy
 - Pregnancy
 - Obesity (BMI > 30) (the 95th percentile)
 - Healthy children under 2 years old are also considered **High-Risk** (Reason: higher rate of pneumonia and hospitalization)
 - Note: All other children are referred to as **Low-Risk** in this guideline (not High-Risk)
88. **Exposure of High-Risk Patient:**
- If the child with flu symptoms lives with a **High-Risk** person (such as a sibling or adult with a chronic disease, persons under 2 years or over 65 years, a pregnant woman, etc), the family needs to call the **High-Risk** patient's PCP within 24 hours. (Reason: may need anti-viral medication.)
89. **Observation During Sleep:**
- Sleep in the same room for a few nights. Reason: Stridor can progress to difficulty breathing and can get worse at night.
90. **Runny Nose - Blow or Suction the Nose:**
- The nasal mucus and discharge is washing viruses and bacteria out of the nose and sinuses.
 - Having your child blow the nose is all that is needed. Teach your child how to blow the nose at age 2 or 3.
 - For younger children, gently suction the nose with a suction bulb. Use saline (salt water) nose drops or spray to loosen up the dried mucus as needed.
91. **How to Protect Others:**
- **Stay Home a Minimum of 10 Days:** Home isolation is needed for at least 10 days after symptoms started.
 - Follow local, state or provincial Department of Health directives.
 - Students should follow their school's COVID-19 policy.
 - Do Not go to stores, restaurants, places of worship or other public places. Avoid public transportation or ride sharing. Do Not allow any visitors (such as friends). Leave the house only if you need to seek medical care.
 - **Avoid Close Contact with Other Family Members:** That includes hugging, kissing, sitting next to or sleeping in the same bed. None of this is realistic for young children. Carefully avoid any contact with the elderly and people with weak immune systems or other chronic health problems.
 - **Wash Hands Often with Soap and Water:** Use an alcohol-based hand sanitizer if water is not available. Don't touch your eyes, nose or mouth unless your hands are clean.
 - **Don't Share Personal Household Items:** Don't share glasses, plates or eating utensils.
 - **Older Children Wear a Mask** in common household areas.

92. **Lab Test Confirmed COVID-19 and NO Symptoms:**
- Your child had a lab test for COVID-19 and it came back positive.
 - A positive result on a PCR or rapid self-test kit is highly accurate for diagnosing COVID-19. It is highly likely that your child has COVID-19.
 - From what you have told me, your child has no symptoms. That is reassuring.
 - Your child will still need home isolation for 10 days after the date of the positive test.
 - Here's some care advice that should help.
93. **Questions about a Negative Rapid COVID-19 Test Result:**
- Positive rapid test results are accurate and can be trusted.
 - Negative rapid test results are usually accurate but can sometimes be wrong.
 - An error is more likely with tests performed at home. Rapid tests performed at a test site are usually more accurate.
 - The COVID-19 vaccine does NOT affect the results of a COVID-19 test.
 - Note to Triager: For callers who are worried about a false negative, especially if they had a known exposure, put the call back to the PCP during normal hours.
 - Tell the caller: "Your doctor will help decide if a special lab test would be useful for your child".
94. **Antibody Tests - Rarely Needed:**
- **Antibody Tests:** These tests are different. These are performed on blood. They can sometimes tell us if there are antibodies from a previous infection. They require a doctor's order and are rarely helpful. If you have questions, discuss with your doctor during office hours.
 - **Timing guideline for Antibody Tests:** If indicated, antibody tests are not recommended until at least 2 or 3 weeks have passed since the start of the infection (CDC). Waiting for a few weeks will give the most accurate result (highest positive rate).
95. **COVID-19 Vaccines - Vaccine Postponement Questions:**
- **Positive COVID-19 Test:** If your child has a positive COVID-19 test, the vaccine should be postponed until after the 10 day quarantine period is over and the symptoms are improving.
 - **Child is Sick and Scheduled for Vaccine:** If your child has symptoms compatible with COVID-19, should get a test before receiving the vaccine. If negative and mild illness (such as isolated runny nose or mild diarrhea), can receive the vaccine. For moderate or severe illness (including a fever), the vaccine should be postponed until fever gone and symptoms are improving.
 - **Exposed to COVID-19, But No Symptoms:** If your child has been exposed to COVID-19 and is scheduled for the vaccine, the vaccine should be postponed until after the quarantine period is over.
 - **Flu and COVID-19 Vaccines:** Can be given at the same time. No waiting period needed between the 2 shots.

FIRST AID



N/A

BACKGROUND INFORMATION

COVID-19 Main Symptoms (CDC)

COVID-19 should be suspected in people who have 1 or more of the following:

- Cough
- Shortness of breath (difficulty breathing)
- Fever or chills
- Loss of smell or taste
- Muscle or body aches
- Headache
- Sore throat
- Runny nose (not from allergies)
- The CDC also includes the following less common symptoms: fatigue (tiredness), nausea, vomiting and diarrhea. In isolation, these symptoms (such as diarrhea) are not very helpful for recognizing COVID-19. Reason: Too common, multiple causes and sometimes subjective. For example, mild diarrhea is often caused by a change in the diet.
- **"COVID Toes"**: Reddish or purple toes have been reported as a rare finding. They can occur alone and go away without treatment. Or they can occur 1-2 weeks after the more common symptoms.

Matching Pediatric Care Advice (PCA) Handouts for Callers

Detailed home care advice instructions have been written for this protocol. If your software contains them, they can be sent to the caller at the end of your call. Here are the names of the pediatric handouts that can be used with this protocol:

- COVID-19 - Diagnosed or Suspected
- COVID Exposure
- COVID-19 Prevention
- COVID-19 or Influenza - How to Tell
- COVID-19 Vaccines - Answers to Common Questions
- Fever - How to Take the Temperature
- Fever - Facts Versus Myths
- Acetaminophen (Tylenol) Dosage Table - Children
- Ibuprofen (Advil, Motrin) Dosage Table - Children

Multisystem Inflammatory Syndrome (MIS-C)

- MIS-C is a very rare complication of COVID-19. In general, COVID-19 continues to be a mild disease in children. It cannot be predicted who will get this complication.
- Prevention: MIS-C can be prevented by getting your child vaccinated against COVID-19.
- The most common symptoms are fever, a red rash, abdominal pain with vomiting and diarrhea. Half of the patients develop trouble breathing. Some children become confused or overly sleepy. Always has multiple symptoms.
- Onset of symptoms: Usually about 4 weeks after a COVID-19 infection and apparent recovery.
- Peak age: 8 years. Age range: 6 months to 21 years.
- Treatment: Most patients with MIS-C need to be admitted to the hospital. MIS-C is treatable with medications, including IV immune serum globulin and steroids.
- Prognosis: Most children with MIS-C have a full recovery. The death rate is about 1 per 100.

Cause

- It is caused by a novel (new) coronavirus (COVID-19).
- Viruses change through mutation. New variants of the COVID-19 virus are expected to appear and spread.
- In the summer and fall of 2021, the Delta variant has become the most common COVID-19 variant.
- The Delta variant spreads much faster than other variants.
- It may cause more severe illness and more hospitalizations.

- The COVID-19 vaccines help protect against the delta variant.
- Infection with COVID-19 Delta variant occurs less often in people who are vaccinated. When it happens it is called a "breakthrough" infection.
- The risk of serious illness and hospitalization is much lower than if a person was not vaccinated.
- Current evidence suggests that vaccinated people who become infected with COVID-19 can also spread the virus to others.

COVID-19 Origins

- An outbreak of this new viral infection began in Wuhan, China in early December 2019.
- The first COVID-19 patient in the United States was reported on January 21, 2020.
- Four patients were confirmed in Canada on January 31, 2020.
- The World Health Organization (WHO) declared COVID-19 a global pandemic on March 11, 2020.
- The Centers for Disease Control and Prevention (CDC) is considered the source of truth for this guideline. This continues to be a rapidly changing situation and recommendations from the CDC are updated daily. See: <https://www.cdc.gov/coronavirus>. If the CDC recommendations are different than what is in this guideline, follow the CDC guidelines.

High-Risk Children for Complications with Influenza (also with COVID-19)

- Significance: HIGH-RISK children also are the main patients who may need prescription anti-viral medications when they develop influenza. New anti-viral medications for COVID-19 may also be indicated for similar patients.
- Lung disease (e.g., asthma, cystic fibrosis, bronchopulmonary dysplasia)
- Technology-dependent lung disease (e.g., oxygen required, tracheostomy, ventilator)
- Compromised ability to handle respiratory secretions (e.g., spinal cord or brain injury)
- Heart disease (e.g., congenital heart disease, rheumatic heart disease)
- Neuromuscular disease (e.g., muscular dystrophy, cerebral palsy, epilepsy)
- Metabolic disease (e.g., diabetes mellitus)
- Sickle cell disease
- Renal disease (e.g., nephrotic syndrome, renal dialysis)
- Liver disease (e.g., liver failure, chronic hepatitis)
- Down syndrome
- Compromised immune system (e.g., cancer, chemotherapy, HIV/AIDS, transplant, taking oral steroids)
- Diseases requiring long-term aspirin therapy (e.g., Kawasaki's disease and rheumatoid arthritis)
- Pregnancy
- Obesity (BMI > 30) (the 95th percentile)
- Healthy children under 2 years old are also considered HIGH-RISK. Reason: higher rate of pneumonia and hospitalization.

High-Risk Children: Possible Exceptions

- The current HIGH-RISK list includes over 20% of children because 10% of children are under 2 years of age and 10% of children have asthma.
- To reduce unnecessary prescribing of Tamiflu, our call center and ED have decided to exclude children who only have exercise-induced asthma or cough-variant asthma. We have also excluded any child with asthma who has not needed to use any asthma medications within the last year. The latter would indicate that they have very mild intermittent asthma.
- Each call center and medical advisory group will need to decide if certain conditions will not be included in the HIGH-RISK group.

Flu without Fever

- Influenza can sometimes occur without fever.
- However, fever remains a valuable aid in triage, and without it the telephone diagnosis of suspected flu could double or triple (because it would include many children with the common cold or very mild cases of influenza). That would overwhelm the health care delivery system.
- Premise: anyone who has flu without a fever has a very mild case of the infection and doesn't need anti-viral medications. (Exception: immune-suppressed individuals who may not be able to mount a febrile response).
- Our call center has decided to exclude children who may have flu respiratory symptoms but do not have a fever. Each call center will need to make their own decision.

Child Abuse During the COVID-19 Pandemic

- The pandemic has increased the incidence of abuse and domestic violence due to social isolation and financial burdens.
- Also, young children often become irritable and demanding when confined to the home.
- Triagers need to be alert for calls about bruises or other injuries that are suspicious, unexplained or occur in the first year of life.
- Offer help to families in crisis before they reach the breaking point. Be alert to increased domestic violence. Know where to refer at-risk families.
- See the Psychosocial Problems, Child Abuse or Domestic Violence guidelines for details.

COVID-19 - Other Facts

- **Incubation Period:** average 5 days (range 2 to 14 days) after coming in contact with the secretions of a person who has COVID-19.
- **No Symptoms but Infected:** Over 30% of infected adult patients have no symptoms (asymptomatic patients). Children and teens are even more likely to have no symptoms. Such patients do however spread the disease and most develop protective antibodies (immunity).
- **Mild Infections:** 80% of adults with symptoms have a mild illness, much like normal flu or a bad cold. The symptoms usually last 2 weeks.
- **Severe Infections:** 20% of unvaccinated adults with symptoms develop trouble breathing from viral pneumonia. Many of these need to be admitted to the hospital. About 2% of unvaccinated children with COVID-19 need to be admitted to the hospital. About 10% of unvaccinated teens need hospitalization. About 3% require ICU care. (CDC). People with complications generally recover in 3 to 6 weeks. Severe infections are rare in people who are vaccinated.
- **Long-Haul Symptoms:** Have been reported in some children after hospitalization with severe infections. Main symptoms are fatigue, brain fog, muscle pains and joint pains. Up to 2% have symptoms beyond 8 weeks.
- **Deaths:** Children generally have a mild illness and recover quickly. Pediatric deaths are very rare. (CDC) Older adults, especially those with chronic lung disease, heart disease, diabetes, obesity or weak immune systems, have the highest death rates. The overall death rate is around 2 per 1000 people. Over 90% of deaths occur in people who are not vaccinated.
- **Vaccine:** Safe and effective vaccines are available. Some vaccines are 2 doses, given 3-4 weeks apart. Others are a single dose. Similar to flu shots, they will probably provide protection for 6 to 9 months. At this time, vaccines have been tested and are FDA approved for 5 years and older. The COVID-19 vaccine will reduce the chance of your child getting COVID-19. The vaccine prevents almost all hospital admissions, ICU care and deaths.
- **"Breakthrough Cases":** These are COVID-19 infections that bypass vaccine protection. They are rare and many are asymptomatic.
- **Treatment:** New treatments for severe COVID-19 are available. They are mainly used on hospitalized patients and are given in a vein (IV). **Caution** - only discuss the following if caller asks about the new anti-viral pill (paxlovid): Paxlovid is given by mouth during the first 3 days of symptoms to prevent serious complications. It awaits FDA approval and will initially be used for adults at high risk for severe disease.
- **Prevention:** The COVID-19 vaccine is the best way to prevent infections. Face masks, social (safe)

distancing and extra handwashing are also proven to help prevent disease. **Caution** - only discuss the following if caller asks about monoclonal antibody therapy: Is available for those 12 and older at high risk for severe disease who are 1) asymptomatic and had a recent close contact exposure OR 2) have confirmed COVID-19 mild symptoms. It is usually given IV to prevent progression and complications. People hospitalized with COVID-19 are not eligible.

COVID-19 - Exposure Risk Factors

- The following are the main risk factors for getting sick with COVID-19:
- **Close contact with a person who tested positive for COVID-19** AND contact occurred while they were ill. Close contact is defined as being within 6 feet (2 meters) for a total of 15 minutes or more over a 24-hour period. Prolonged close contact would extend the risk to the 48 hours prior to the person becoming ill with symptoms. This includes **living with someone infected** with COVID-19.
- **Living in or travel to an area** where there is **high community spread** of COVID-19 also carries some risk.
- **International travel:** The CDC (<https://www.cdc.gov/coronavirus>) has the most up-to-date list of where COVID-19 outbreaks are highest.
- **Not being fully vaccinated**

Animals and COVID-19

- The main way COVID-19 spreads is from person to person. There is low risk of getting COVID-19 from a pet or other animal.
- It is possible for animals to catch COVID-19 from people. A few pets have tested positive for COVID-19 (including cats and dogs).
- The CDC recommends treating pets like other family members when trying to avoid spreading COVID-19. Do not let pets have close contact with other people or animals outside your household. A sick person should self-isolate and avoid contact with both people and pets.
- Call your vet if your pet gets sick or you have other questions.
- The CDC has more information on COVID-19 and animals at: <https://www.cdc.gov/coronavirus>.

COVID-19 Disease and Repeat Infections

- Most viral infections cause our immune system to create antibodies that protect us from getting that infection again.
- Sometimes this provides lifelong protection, but sometimes that protection only lasts months or years.
- **Protection Duration after an Infection.** Research about how long protection against COVID-19 lasts is ongoing. Protection has been proven to last for at least 90 days (3 months) after infection. The CDC recommends using 90 days post exposure as a protected period.
- For now, it remains important for people who have recovered from COVID-19 infections to be careful. Take normal precautions such as wearing a mask and social distancing.
- **Need for Vaccine.** People who have recovered from COVID-19 should still get a COVID-19 vaccine. Reason: Vaccination provides greater protection than the natural immunity from a COVID-19 infection. The greatest protection comes from having both (CDC).
- **Recovery and Re-infections.** Re-infections after full recovery do occur. The arrival of COVID-19 variant (mutant) viruses has increased the rate of re-infections for some of the variants.
- **Vaccines and Break-through Infections.** COVID-19 vaccines protect against most of the COVID-19 variants. Even when they don't, they usually protect against severe disease and the need for hospitalization.
- **Booster Vaccines:** Booster vaccines are recommended 6 or more months after the Pfizer or Moderna vaccines and 2 or more months after the Johnson and Johnson vaccine. These booster shots reduce the rate of COVID-19 break-through infections. Initially, they are recommended for people at high risk of severe disease. Soon they will be available to everyone.

Ibuprofen and other NSAID Use for COVID-19

- Many callers have expressed concerns that ibuprofen (or other NSAID) use to treat COVID-19 symptoms may worsen the disease.
- These concerns originated from a few physicians' comments and have since spread over social media.
- To date, there is no scientific evidence (clinical trials or studies) that show that using ibuprofen negatively impacts outcome in COVID-19 patients. We will continue to review any new literature as it is published.
- The CDC, WHO, AAP and our Infectious Disease expert reviewers continue to approve the use of ibuprofen for COVID-19.
- For this reason, STCC guidelines continue to recommend ibuprofen as an acceptable way to treat high fevers and pain. (Note: Remind callers that fevers are beneficial, help fight the infection, and may speed recovery. Low-grade fevers should not be treated.)
- If callers remain concerned, they can use acetaminophen for symptoms that warrant treatment.
- Caution: For suspected COVID-19 patients on oral steroids, such as prednisone, the triager should involve the PCP for a decision about whether the drug can be continued.

Internet Resources

- Centers for Disease Control and Prevention (CDC): Coronavirus. <https://www.cdc.gov/coronavirus>.
- Public Health Agency of Canada: <https://www.canada.ca/en/public-health/services/diseases/coronavirus.html>.
- World Health Organization (WHO): Coronavirus. <https://www.who.int/health-topics/coronavirus>.
- American Academy of Pediatrics: <http://www.healthychildren.org>

Expert Reviewers

- Jessica Cataldi, MD, Pediatric Infectious Diseases and Epidemiology, Children's Hospital Colorado, Aurora, Colorado
- Samuel Dominguez, MD, Pediatric Infectious Diseases and Epidemiology, Children's Hospital Colorado, Aurora, Colorado
- Lisa M. Koonin DrPH, MN, MPH; Founder, Health Preparedness Partners; Pandemic preparedness specialist
- The author is extremely grateful for these critical reviews.

REFERENCES

1. Alramthan A, Aldaraji W. A case of COVID-19 presenting in clinical picture resembling chilblains disease. First report from the Middle East. Clin Exp Dermatol 2020 Aug;45(6):746-748.
2. Bautista-Rodriguez C, Sanchez-de-Toledo J, Clark BC, et al. Multisystem Inflammatory Syndrome in children: An international survey. Pediatrics 2021 Feb;147(2):e2020024554.
3. Castagnoli R, Votto M, Licari A, et al. Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Infection in Children and Adolescents: A Systematic Review. JAMA Pediatr. 2020 Sep 1;174(9):882-889.
4. CDC COVID-19 Response Team. Coronavirus Disease 2019 in Children - United States, February 12 - April 2, 2020. MMWR Morbidity and Mortality Weekly Report. ePub: 6 April 2020.
5. Chung E, Chow EJ, Wilcox NC, et al. Comparison of Symptoms and RNA Levels in Children and Adults With SARS-CoV-2 Infection in the Community Setting. JAMA Pediatr. 2021 Jun 11.

6. De Rose DU, Piersigilli F, Ronchetti MP, et al. Novel coronavirus (COVID-19) in newborns and infants. *Ital J Pediatr*. 2020 Apr 29;46(1):56.
7. DeLaroche AM, Rodean J, Aronson PL, et al. Pediatric Emergency Department visits at US Children's Hospitals during the COVID-19 pandemic. *Pediatrics*. 2021 Apr;147(4):e2020039628.
8. Dufort EM, Koumans EH, Chow EJ, et al. Multisystem Inflammatory Syndrome in children in New York state. *N Engl J Med*. [published online ahead of print, 2020 Jun 29]
9. Farooqi KM, Chan A, Weller RJ, et al. Longitudinal Outcomes for Multisystem Inflammatory Syndrome in Children. *Pediatrics*. 2021 Aug;148(2):e2021051155.
10. Feldstein LR, Rose EB, Horwitz SM, et al. Multisystem Inflammatory Syndrome in U.S. children and adolescents. *N Engl J Med*. [published online ahead of print, 2020 Jun 29].
11. Fernandes DM, Oliveira CR, Guerguis S, et al. Severe Acute Respiratory Syndrome Coronavirus 2 Clinical Syndromes and Predictors of Disease Severity in Hospitalized Children and Youth. *J Pediatr*. 2021 Mar;230:23-31.e10.
12. Fouda GGA, Kwiek JJ, Yotebieng M. Safety of breastfeeding by mothers with COVID-19: New evidence from Israel. *Pediatrics*. 2021 Apr 13;e2020049772.
13. Harrison E, Garbutt J, Sterkel R, et al. Collaborating to advocate in primary care for children during COVID-19. *Pediatrics*. 2021 Oct;148(4):e2021052106.
14. Hatoun J, Correa ET, Donahue SMA, et al. Social distancing for COVID-19 and diagnoses of other infectious diseases in children. *Pediatrics*. 2020 Oct;146(4):e2020006460.
15. Hernandez C, Bruckner AL. Focus on "COVID Toes". *JAMA Dermatol*. 2020 Sep 1;156(9):1003.
16. Humphreys KL, Myint MT, Zeanah CH. Increased risk for family violence during the COVID-19 pandemic. *Pediatrics*. 2020 Jul;146(1):e20200982.
17. Jain SS, Steele JM, Fonseca B, et al. COVID-19 Vaccination - Associated Myocarditis in Adolescents. *Pediatrics*, Nov 2021, 148 (5) e2021053427.
18. Kainth MK, Goenka PK, Williamson KA, et al. Early experience of COVID-19 in a US children's hospital. *Pediatrics*. 2020 Oct;146(4):e2020003186.
19. King JA, Whitten TA, Bakal JA, et al. Symptoms associated with a positive result for a swab for SARS-CoV-2 infection among children in Alberta. *CMAJ*. 2021 Jan 4;193(1):E1-E9.
20. Laws RL, Chancey RJ, Rabold EM, et al. Symptoms and transmission of SARS-CoV-2 among children - Utah and Wisconsin, March-May 2020. *Pediatrics*. 2021 Jan;147(1):e2020027268.
21. Lu X, Zhang L, Hui, D, et al. SARS-CoV-2 Infection in children. *N Engl J Med*. 2020 Apr 23;382(17):1663-1665.
22. Ludvigsson JF. Systematic review of COVID-19 in children shows milder cases and a better prognosis than adults. *Acta paediatrica*. March 2020. doi:10.1111/apa.15270.
23. McCormick DW, Richardson LC, Young PR, et al. Deaths in Children and Adolescents Associated With COVID-19 and MIS-C in the United States. *Pediatrics*, Nov 2021, 148 (5) e2021052273.
24. Mithal LB, Machut KZ, Muller WJ, et al. SARS-CoV-2 infection in infants less than 90 days old. *J Pediatr* 2020 Sep;224:150-152.

25. Muchmore B, Muchmore P, Lee CW, et al. Tracking potential COVID-19 outbreaks with influenzalike symptoms urgent care visits. *Pediatrics*. 2020 Oct;146(4):e20201798.
26. Ouldali N, Yang DD, Madhi F, et al. Factors associated with severe SARS-CoV-2 infection. *Pediatrics* March 2021,147 (3) e2020023432.
27. Paret M, Lalani K, Hedari C, et al. SARS-CoV-2 among infants <90 days of age admitted for serious bacterial infection evaluation. *Pediatrics*. 2021 Oct;148(4):e2020044685.
28. Romero Ramírez DS, Lara Pérez MM, Carretero Pérez M, et al. SARS-CoV-2 Antibodies in Breast Milk After Vaccination. *Pediatrics*, Nov 2021, 148 (5) e2021052286.
29. Ruiyun Li, Sen Pei, Bin Chen, et al. Substantial undocumented infection facilitates the rapid dissemination of novel coronavirus (SARS-CoV2). *Science* 10.1126/science.abb3221 (2020)
30. Shekerdemian LS, Mahmood NR, Wolfe KK, et al. Characteristics and outcomes of children With Coronavirus Disease 2019 (COVID-19) infection admitted to US and Canadian pediatric intensive care units. *JAMA Pediatr*.2020 Sep 1;174(9):868-873.
31. Shlomai NO, Kasirer Y, Strauss T, et al. Neonatal SARS-CoV-2 infections in breastfeeding mothers. *Pediatrics*. 2021 May;147(5):e2020010918.
32. Song W, Li J, Zou N, et al. Clinical features of pediatric patients with coronavirus disease (COVID-19). *J Clin Virol*. 2020 Apr 24;127:104377.
33. Su L, Ma X, Yu H, et al. The different clinical characteristics of corona virus disease cases between children and their families in China - the character of children with COVID-19. *Emerging Microbes and Infection* 2020; 9(1): 707-13.
34. Szilagyi PG, Shah MD, Delgado JR, et al. Parents' Intentions and Perceptions About COVID-19 Vaccination for Their Children: Results From a National Survey. *Pediatrics*. 2021 Oct;148(4):e2021052335.
35. Wong CA, Ming D, Maslow G, et al. Mitigating the impacts of the COVID-19 pandemic response on at-risk children. *Pediatrics*. 2020 Jul;146(1):e20200973.
36. Zimmerman KO, Brookhart MA, Kalu IC, et al. Community SARS-CoV-2 Surge and Within-School Transmission. *Pediatrics*. 2021 Oct;148(4):e2021052686.

SEARCH WORDS

2019-NCOV
 BREATHING
 BREATHING DIFFICULTY
 CORONAVIRUS
 CORONAVIRUS EXPOSURE
 COUGH
 COVID
 COVID TOES
 COVID-19
 DIFFICULT BREATHING
 DIFFICULTY BREATHING
 EXPOSURE
 EXPOSURE QUESTION

EXPOSURE QUESTIONS
FOREIGN TRAVEL
INFECTION EXPOSURE
INTERNATIONAL TRAVEL
MIS-C
MULTISYSTEM INFLAMMATORY SYNDROME
NCOV
NOVEL CORONAVIRUS
SARS-COV-2
SOB
TRAVEL
TROUBLE BREATHING

AUTHOR AND COPYRIGHT

Author: Barton D. Schmitt, MD, FAAP
Copyright: 1994-2021, Schmitt Pediatric Guidelines LLC All rights reserved.
Company: Schmitt-Thompson Clinical Content
Content Set: After Hours Telehealth Triage Guidelines | Pediatric
Version Year: 2021
Last Revised: 11/13/2021
Last Reviewed: 11/13/2021